

Viroids: Survivors from the RNA World?

Ricardo Flores,¹ Selma Gago-Zachert,²
Pedro Serra,¹ Rafael Sanjuán,³ and Santiago F. Elena^{1,4}

¹Instituto de Biología Molecular y Celular de Plantas (UPV-CSIC), 46022 València, Spain; email: rflores@ibmcp.upv.es

²Abteilung Molekulare Signalverarbeitung, Leibniz Institut für Pflanzenbiochemie, 06120 Halle (Saale), Germany

³Instituto Cavanilles de Biodiversidad y Biología Evolutiva, Universidad de València, 46980 València, Spain

⁴The Santa Fe Institute, Santa Fe, New Mexico 87501

Annu. Rev. Microbiol. 2014. 68:395–414

First published online as a Review in Advance on June 18, 2014

The *Annual Review of Microbiology* is online at micro.annualreviews.org

This article's doi:
10.1146/annurev-micro-091313-103416

Copyright © 2014 by Annual Reviews.
All rights reserved

Keywords

viroid-like satellite RNAs, hepatitis δ virus, catalytic RNAs, hammerhead ribozymes, small noncoding RNAs

Abstract

Because RNA can be a carrier of genetic information and a biocatalyst, there is a consensus that it emerged before DNA and proteins, which eventually assumed these roles and relegated RNA to intermediate functions. If such a scenario—the so-called RNA world—existed, we might hope to find its relics in our present world. The properties of viroids that make them candidates for being survivors of the RNA world include those expected for primitive RNA replicons: (*a*) small size imposed by error-prone replication, (*b*) high G + C content to increase replication fidelity, (*c*) circular structure for assuring complete replication without genomic tags, (*d*) structural periodicity for modular assembly into enlarged genomes, (*e*) lack of protein-coding ability consistent with a ribosome-free habitat, and (*f*) replication mediated in some by ribozymes, the fingerprint of the RNA world. With the advent of DNA and proteins, those protoviroids lost some abilities and became the plant parasites we now know.

Contents

INTRODUCTION: WHY THE NEED FOR AN RNA WORLD?	396
VIROIDS: ESSENTIAL FEATURES	397
Discovery of a Subviral World	397
Structure: Small Circular RNAs with Compact Folding	397
Replication: Rolling-Circle Mechanism Catalyzed by Enzymes and Ribozymes	399
Sequence Diversity	400
VIROID-RELATED REPLICONS	401
Viroid-Like Satellite RNAs	401
Retroviroid-Like Elements	401
<i>Hepatitis δ virus</i>	402
WHY ARE VIROIDS AND VIROID-RELATED REPLICONS REGARDED AS	
SURVIVORS OF THE RNA WORLD?	402
Early Speculations on the Origin of Viroids	402
“Circular RNAs: Relics of Precellular Evolution?”	402
Transition from the RNA World to a World with Proteins and DNA: Viroids	405
Transition from the RNA World to a World with Proteins and DNA:	
Viroid-Related Replicons	407
SUMMARY	407

INTRODUCTION: WHY THE NEED FOR AN RNA WORLD?

The role of RNA has long been considered a subsidiary one, a mere link between DNA and proteins, which store and express genetic information in all cells. However, this is not true in the subcellular world, because the genetic material of many viruses—including that of *Tobacco mosaic virus* (TMV), the first discovered (4)—is RNA (48, 55). Although replication of some RNA viruses infecting bacteria (bacteriophages) was initially deemed to proceed through a mechanism different from that proposed for DNA (68), ensuing work with bacteriophage Q β demonstrated that its single-stranded RNA (ssRNA) of about 4,200 nucleotides (nt) replicates by synthesizing a complementary strand via base pairing between A and U (instead of A and T, as in DNA), and G and C (41). Therefore, in these very simple (and presumably very old) self-replicating entities, RNA supplants DNA as the carrier of genetic information. Subsequent discovery of ribozymes, i.e., that some RNAs have catalytic activity (61, 81), showed that RNA can also express this information, supporting an early proposal that RNA could be the primordial macromolecule (18, 101, 143). This hypothesis was based on the ability of ssRNA to fold into complex structures—hairpins with stems stabilized by hydrogen bonds between bases and with loops formed by bases with no partners available—resembling those of proteins (50), and on its 2' hydroxyl groups providing additional chemical versatility.

The term RNA world was coined in a commentary (56) following discovery of ribozymes:

If there are ribozymes that can catalyze the synthesis of a new RNA molecule from precursors and an RNA template, then there is no need for protein at the beginning of evolution. One can contemplate an RNA world, containing only RNA molecules that serve to catalyze the synthesis of themselves. . . . At the next stage RNA molecules began to synthesize proteins, first by developing RNA adapter molecules that can bind activated amino acids, and then by arranging them according to an RNA template using other

RNA molecules such as the RNA core of the ribosome. This process would make the first proteins, which would simply be better enzymes than their RNA counterparts. . . . Finally, DNA appeared on the scene, the ultimate holder of information copied from the genetic RNA molecules by reverse transcription. After double-stranded DNA (dsDNA) evolved there exists a stable linear information store, error correcting because of its double-stranded structure but still capable of mutation and recombination. RNA is then relegated to the intermediary role that it has today—no longer the center of the stage, displaced by DNA and the more efficient protein enzymes.

This enlightening proposal has been extended into an entire field of research (3). Implicit in the RNA world idea is the existence of primitive RNA replicons of minimal size that later evolved into more complex self-replicating systems. Are there remnants of those minimal replicons in our present-day world? In the following sections we provide evidence that viroids (and some viroid-related replicons) are excellent candidates for being survivors from the RNA world. But first we summarize the unique, pertinent properties of these genetic elements. In-depth reviews dealing with different aspects of viroids have been published previously (30–32, 42, 43, 45, 66, 95, 128, 132).

VIROIDS: ESSENTIAL FEATURES

Discovery of a Subviral World

A recurrent situation in science is that unanticipated discoveries entail unanticipated consequences going beyond the initial scheme. Viroids were discovered during an attempt to identify the agent of potato spindle tuber (PST) disease, suspected to be a virus. The finding that the PST agent can be transmitted to tomato, which is easier to grow and expresses symptoms more rapidly, expedited the experiments and produced the first surprise: After ultracentrifugation of tomato extracts, the infectious principle remained in the supernatant instead of being associated with the presumed viral particles—the virions, characteristically formed by protein subunits encapsidating the nucleic acid—expected to accumulate in the sediment. Besides, this infectious principle moved in sucrose gradient centrifugation more slowly than typical virions, and its mobility remained unchanged after deproteinating treatments, indicating that the PST agent might be a naked nucleic acid. This hypothesis was tested with polyacrylamide gel electrophoresis (PAGE): The PST infectious principle migrated as a band, absent in mock-inoculated plants, with the mobility predicted for a minuscule RNA (because of its sensitivity to RNase, but not to DNase). At this stage the term viroid was proposed for what appeared to be the first autonomously replicating subviral agent (*Potato spindle tuber viroid*, PSTVd) (26, 27). Application of this methodology quickly resulted in discovery of novel viroids as the causal agents of other plant diseases of presumed, but not proved, viral etiology. Altogether these studies opened a window onto a subviral world composed of small, self-replicating RNAs, displacing viruses from the lowest step of the biological scale, which they had occupied during the first 65–70 years of the twentieth century. It is worth noting that the subviral world, like the viral world, was discovered by studies on plant systems. However, in contrast to viruses, viroids so far have been restricted to the plant kingdom, although at least one viroid-related replicon is involved in a human disease (see below).

Structure: Small Circular RNAs with Compact Folding

First of all, how small was PSTVd compared with the RNAs of known viruses? Early estimations from PAGE, corroborated by electron microscopy of purified PSTVd RNA (124), were about 300 nt, a size at least 10-fold smaller than that of the ssRNAs of TMV (6,400 nt) and bacteriophage

Q β (see above). Moreover, electron microscopy of denatured PSTVd RNA produced a second surprise: The RNA was circular, a feature previously conjectured given its resistance to exonucleases. Together, the small size (the RNA encoding, if any, translation products of minimal complexity) and circularity (which hampered translation by conventional ribosome scanning from a 5' terminus) suggested that PSTVd was a nonprotein-coding RNA. Determination of its 359-nt primary structure by direct RNA sequencing (a milestone in molecular biology, given that it was the first eukaryotic pathogen completely sequenced) confirmed the circular structure and the absence of typical initiation codons (59). From a functional perspective this was a distinctive point, because all viruses encode proteins mediating one or more steps of their biological cycle. Furthermore, thermodynamic predictions (109), along with RNase and bisulfite probing *in vitro*, supported the idea of intramolecular folding of PSTVd into a rodlike secondary structure—also favored by circularity and composed of double-stranded segments flanked by apparently unstructured loops—with a width akin to that of a dsDNA as observed by electron microscopy under nondenaturing conditions.

Sequencing of other viroids confirmed preservation of the rodlike folding and, within it, of a central conserved region (CCR) (90), which emerged as an idiosyncratic trait (**Figure 1**). However, this was not the case of *Avocado sunblotch viroid* (ASBVd) (126), which did not have a CCR but had strands of both polarities displaying a remarkable behavior: They self-cleaved through




Figure 1

Viroid structure. (*a*) Rodlike secondary structure with a central conserved region (CCR) proposed for PSTVd (family *Pospiviroidae*). (*b*) Branched secondary structures proposed for PLMVd (family *Avsunviroidae*) and (*c*) for its plus-polarity hammerhead ribozyme. Nucleotides strictly or highly conserved in natural hammerhead structures are shown within boxes with orange and blue backgrounds for plus and minus polarities, respectively. Continuous lines and dots between nucleotides represent canonical (Watson-Crick) and noncanonical base pairs, respectively, with the red arrow marking the self-cleavage site. Dashed rectangles denote tertiary interactions between loops that either stabilize the global viroid conformation or promote the catalytically active ribozyme folding.




Figure 2

Viroid classification. Distinctive structural and functional features of the two viroid families, *Pospiviroidae* (from POtato SPindle tuber VIROID) and *Avsunviroidae* (from AVocado SUNblotch VIROID).

hammerhead ribozymes, thus acting as catalytic RNAs (74), an attribute with deep implications for its replication and evolutionary origin. PSTVd and ASBVd epitomized two viroid groups, which ultimately resulted in recognition of the families *Pospiviroidae* and *Avsunviroidae*, respectively (37, 43), a cataloging consistent with other distinctive features that include the subcellular site and mode of replication (see below) (**Figure 2**). Moreover, some members of the *Avsunviroidae*, like *Peach latent mosaic viroid* (PLMVd) (71) and *Chrysanthemum chlorotic mottle viroid* (CChMVd) (93), fold into branched secondary structures (**Figure 1**).

Replication: Rolling-Circle Mechanism Catalyzed by Enzymes and Ribozymes

Because viroids are nonprotein-coding RNAs, they need to usurp the transcription and processing machinery of their hosts to replicate. Thus, viroids are essentially transcriptional parasites, whereas viruses are essentially translational parasites (to infect hosts, they must express the proteins they encode). Viroid replication takes place in the nucleus (family *Pospiviroidae*) or in plastids, mostly chloroplasts (family *Avsunviroidae*), through a rolling-circle mechanism with RNA intermediates (7, 21, 58, 73). The infecting circular (+) RNA (the sign is conventionally allocated to the most abundant strand *in vivo*) is repeatedly transcribed into oligomeric (–) strands. These replicative intermediates, by themselves (asymmetric pathway) or after being cleaved and ligated into circular monomers (symmetric pathway), serve as templates for synthesis of oligomeric (+) strands that are finally processed into their monomeric circular counterparts (**Figure 3**).

Therefore, three enzymes (or ribozymes) are required: RNA polymerase, RNase, and RNA ligase, the characterization of which has led to unexpected discoveries. First, despite the existence of several RNA-dependent RNA polymerases in plants (118, 138), transcription of viroid strands is catalyzed by nuclear or plastidic DNA-dependent RNA polymerases forced to accept RNA templates (44, 92, 97, 113, 119). Second, cleavage of oligomeric strands is mediated in the family *Pospiviroidae* by a class III RNase enzyme that operates on a dsRNA structure formed by the upper CCR strand of two consecutive units (53), or by *cis*-acting hammerhead ribozymes in the family *Avsunviroidae* (74). And third, circularization occurs by a plastidic isoform of the tRNA ligase (99) or, remarkably, by nuclear DNA ligase 1 (redirected to circularize RNA substrates) (98). Therefore, manifesting their extraordinary parasitic abilities, viroids reprogram the template specificity of a DNA-dependent RNA polymerase to function as an RNA-dependent RNA polymerase, and the substrate specificity of a DNA ligase to operate as an RNA ligase. To complete their infectious




Figure 3

Asymmetric and symmetric variants of the rolling-circle mechanism proposed for replication of members of the families *Pospiviroidae* and *Avsunviroidae*, respectively. Orange and blue colors refer to plus and minus polarities, respectively, with cleavage sites denoted by arrowheads. The enzymes and ribozymes that presumably catalyze the replication steps are indicated. Notice that RNA polymerase II (and NEP) is redirected to transcribe RNA templates and DNA ligase 1 to circularize RNA substrates. Abbreviations: HHRz, hammerhead ribozyme; NEP, nuclear-encoded polymerase.

cycle, viroids also need to move to invade distal plant parts. Recent data indicate that specific loops/bulges in the rodlike structure of PSTVd and related viroids are, instead of unstructured, stabilized by arrays of noncanonical pairs, with some of these elements functioning in replication and others in systemic trafficking (31, 32, 147).

Sequence Diversity

As a consequence of their high mutation rates (see below), some viroids display high genetic variability. This phenomenon has been observed after inoculating plants with infectious viroid cDNAs (usually head-to-tail dimeric constructs) or their transcripts (mimicking replicative intermediates) and quantifying the heterogeneity of the resulting progeny. The diversity observed in the family *Pospiviroidae*, e.g., in PSTVd (57), *Citrus exocortis viroid* (5, 64, 135), *Citrus bent leaf viroid* (52), *Chrysanthemum stunt viroid* (CSVd) (16), and *Citrus dwarfing viroid* (131), is relatively low, and the consensus sequence does not deviate significantly from the wild type. However, in the family *Avsunviroidae*, e.g., PLMVd (2, 94) and CChMVd (16, 23), the resulting population is a complex spectrum of mutants and the consensus sequence often changes over time or is difficult to define. This difference is most likely caused by the distinct replication fidelity of the RNA polymerases involved (see below), with selection imposed by the host also shaping the final distribution of genetic variants (5, 122, 131, 139).

Given their small genomes and high mutation rates, viroids have often been treated in the framework of the quasi-species theory, originally developed to describe natural selection acting on primitive RNA replicons undergoing error-prone replication and consisting of mutant distributions (mutant spectra or clouds) (6, 34). This theoretical scheme was first applied to

bacteriophage QB (33) and then to many other RNA viruses (84, 100, 134). A unique prediction of the quasi-species theory that distinguishes it from classic population genetics models is that selection operates on sets of variants from a particular region of the fitness landscape rather than on individual virus variants. As a result, the fitness associated with a specific sequence depends on the average fitness of its neighbors in the sequence space (84), as first experimentally shown with the RNA bacteriophage $\phi 6$ (11). This collective behavior can lead to the so-called survival of the flattest effect, whereby a population located in a region of the fitness landscape where a large proportion of sequence neighbors are selectively neutral can outcompete another population located in a higher-fitness peak but with a more deleterious sequence neighborhood (121, 141). This effect was first demonstrated *in vivo* by competition experiments between CSVd and CChMVd, performed under different conditions: At the physiological mutation rate the fittest CSVd outcompeted CChMVd, but increasing mutation rate by UV irradiation reversed the outcome and the flattest CChMVd outcompeted CSVd (16). Later, the survival of the flattest was also shown in *Vesicular stomatitis virus* populations subjected to increased mutational stress by chemical mutagenesis (115).

Recent work has suggested that interactions such as genetic complementation and interference among individual components of the viral population also determine to a good extent its behavior, with interference most likely being exerted by defective gene products resulting in nonfunctional protein complexes during the virus biological cycle (100). However, since viroids are nonprotein-coding RNAs, complementation by protein products acting *in trans* cannot exist, and interference must occur at the RNA level.

VIROID-RELATED REPLICONS

Viroid-Like Satellite RNAs

Besides viroids, three other viroid-related subviral elements deserve mention in the context of the RNA world. Plant viroid-like satellite RNAs are structurally similar to viroids, comprising only a small nonprotein-coding circular RNA (114, 120, 127). Additional similarities include replication through a rolling-circle mechanism with ribozyme-mediated self-cleavage of the resulting RNA oligomeric strands of one or both polarities. However, ribozymes are not only of the hammerhead class (46, 106), but also of the hairpin class (12). More important, replication of viroid-like satellite RNAs is not autonomous but relies on the RNA replicase complex encoded—at least in part—by a coinfecting helper RNA virus (whose coat protein encapsidates the satellite RNA for transmission); the term satellite refers to this dual dependence (10). Because the helper RNA viruses replicate in organelle-associated membranous vesicles connected with the cytoplasm (25), replication of viroid-like satellite RNAs most likely also takes place in these vesicles, although lack of direct supporting data leaves open the possibility of the helper virus encoding protein(s) that redirect(s) a host RNA polymerase (127). Resembling the situation found in the family *Avsunviroidae*, circularization of unit-length strands resulting from self-cleavage in viroid-like satellite RNAs might be mediated by a cytoplasmic isoform of the tRNA ligase, or by self-ligation (the RNA ligase activity of the hairpin ribozyme is greater than that of the hammerhead ribozyme) (12, 40).

Retroviroid-Like Elements

A second class of plant small circular RNA is similar to viroid and viroid-like RNAs in size and in harboring hammerhead ribozymes (70) but has a singular property: The RNA is accompanied by a homologous DNA counterpart fused to sequences of either a pararetrovirus or the host (19, 69). This class, dubbed retroviroid-like elements because the homologous DNA is presumably

generated by a reverse transcriptase (RT)—pararetroviruses encode such an enzyme—is restricted so far to some carnation sources, and it cannot be transmitted horizontally. The presence of hammerhead ribozymes in both polarity strands of this carnation circular RNA is evidence of its replication through a symmetric rolling-circle mechanism, but the lack of horizontal transmission suggests that a supply of transcripts from the DNA form is needed.

Hepatitis δ virus

In contrast with the other two, the third class of viroid-related subviral entities has been reported in animals, specifically in humans. With an approximate size of 1,680 nt, the *Hepatitis δ virus* (HDV) RNA—the smallest genome of an animal virus—occupies the position immediately above viroids on the biological scale of genome sizes, and it displays striking structural similarities with viroids: circularity and folding into a rodlike secondary structure (15, 110, 111, 136). Moreover, HDV RNA replicates autonomously in the nucleus through a rolling-circle mechanism catalyzed by host enzymes (an RNA polymerase redirected to transcribe RNA templates and, presumably, an RNA ligase) and by *cis*-acting ribozymes idiosyncratic to this infectious agent (82, 83). However, HDV RNA encodes in its antigenomic polarity a protein (like some RNA viruses) (136), and for transmission it depends on a helper virus, *Hepatitis B virus* (similar to viroid-like satellite RNAs).

WHY ARE VIROIDS AND VIROID-RELATED REPLICONS REGARDED AS SURVIVORS OF THE RNA WORLD?

Early Speculations on the Origin of Viroids

Initial ideas suggested that viroids could be escaped introns, thus descending from host RNAs. This proposal was based on (a) sequence similarities of PSTVd (and other viroids) with group I and II introns, and (b) specific cleavage of viroid oligomeric replication intermediates and their following ligation into circular forms, conceptually resembling cleavage-ligation by which introns are self-spliced (and circularized) from primary transcripts and exons are joined into mature RNAs (28, 63). Yet, subsequent findings that reaction mechanisms were distinct eroded the credibility of the viroid-intron connection.

Moreover, comparative sequence analysis of PSTVd and two related viroids revealed conspicuous similarities with the ends of transposable elements, which together with the presence of inverted repeats and flanking imperfect direct repeats suggested that viroids might have originated from transposable elements or retroviral proviruses by deletion of internal fragments (79). However, despite initial reports indicating the presence of PSTVd sequences in cellular DNA of normal plants, further analyses failed to confirm these results, and moreover, ASBVd (and later other members of its family) did not fit this model. Therefore, viroids do not seem to be simplified versions of viruses or transposons.

“Circular RNAs: Relics of Precellular Evolution?”

With this provoking title, Diener (29) gave a twist to previous perspectives and proposed that viroids and viroid-like satellite RNAs were more likely candidates than introns as living fossils of a precellular RNA world—precellular should be understood to mean lacking cells as they are known today—and he showed that their characteristic features might have evolved to cope with barriers in the self-replication of primitive RNAs. Further studies have reinforced this view and

VIROIDS: THE OLDEST RNA REPLICONS?

Characteristics of viroids giving evidence that they may have a very old origin:

- Minimal genomes (250–400 nucleotides)
- High G + C content, increasing replication fidelity of primitive RNA polymerases
- Circular structure, excluding need of genomic tags for complete replication
- Structural periodicities facilitating modular assembly
- No protein-coding capacity, suggesting emergence before the ribosome
- Presence of ribozymes, the signature of the RNA world

filled some gaps. Hereafter we discuss the evidence that supports Diener's claim, summarizing his arguments and supplying others (see sidebar, Viroids: The Oldest RNA Replicons?).

First, the intrinsic error-prone replication of primitive RNA systems imposed a limit to the size of their master (predominant) sequences to overcome the “error catastrophe” leading to extinction (34). Thus, the evolutionary paradigm that the simplest is the oldest, which with some limitations is generally accepted, seems appropriate for viroids because they are the smallest known replicons.

Second, to increase replication fidelity, (G + C)-rich sequences would have been selected in primitive RNA systems because of the greater thermodynamic stability of GC pairs versus their AU counterparts (35). Viroids also meet this requirement: they are (G + C)-rich with a notable exception, ASBVd (126). Recently, it was proposed that the conflict intrinsic to the RNA world between stable folding (required for ribozyme activity) and template ability (required for replication) could have been solved by a division of labor between the two RNA strands, with one (containing wobble G:U pairs) folding into a ribozyme and the reverse complement (containing the less stable C:A mispairs) functioning as genome (75). Analysis of 40 viroid sequences is consistent with this proposal, a remarkable finding because they must have evolved in response to different selection pressures beyond a simplified division of labor (75). However, this proposal should be restricted to members of the family *Avsunviroidae*, the only ones with hammerhead ribozymes. Furthermore, given that the ribozymes can be formed by the two strands and both function as templates (**Figure 3**), the division of labor view becomes blurred.

Third, circularity allows complete replication without recurring to genomic tags, which mark specific initiation sites in linear templates and ensure their end-to-end copying. Even if polymerization of viroid strands in the present cellular habitat starts at defined sites (24, 80, 96), this may not have occurred in primitive replicons, for which reiterative copying of circular genomes—irrespective of initiation at a defined position—would avoid loss of genetic information. Moreover, this rolling-circle replication would result in multiple copies of the genetic information, a condition likely favored in primitive replicons with a high mutation rate (107). Inherent to this replication mode is the need for a processing mechanism (cleavage and circularization) of the oligomeric replication intermediates (**Figure 3**). Members of the family *Avsunviroidae* fulfill the first step of this mechanism, because their oligomeric strands self-cleave through hammerhead ribozymes; the second step also appears feasible, as revealed by the high level of RNA ligase activity associated with the hairpin ribozymes present in some viroid-like satellite RNAs (10, 12).

Fourth, the sequence of some viroids exhibits a structural periodicity characterized by repeat units of different length. Although this periodicity was initially associated with the protein-binding

ability of viroids (78), other interpretations pointed to a mechanism by which larger genomes could have evolved (29). Even if the strength of this argument is weakened by the lack of structural periodicity in at least one viroid and in most viroid-like satellite RNAs (29), the idea that viroids could have been built modularly seems attractive, particularly if several instead of a single module are considered. The recombinant nature of some viroids, which are chimeras formed by fragments present in other viroids, is consistent with this view (67, 108). Moreover, modular evolution is more advantageous than direct evolution of large functional molecules because of the higher mutation rates allowed, the shortening of evolutionary times, and the possible emergence of complex structures that could not be otherwise directly selected (88). According to computational predictions the module repertoire would be dominated by hairpin-like structures, a fraction of which could have RNA ligase activity and catalyze assembly of larger, eventually functional RNAs; this stepwise ligation-based model of modular evolution could ease the path to the appearance of a ribozyme with RNA replicase activity (9).

Fifth, viroids are nonprotein-coding RNAs. This key feature not only distinguishes viruses from viroids but also traces their putative origin back to what has been called the first age of the RNA world. This first age was delimited between the emergence of the first replicating RNA cell(s)—a metabolism without cellular confinement seems unlikely—and just before the advent of the ribosome, which opened the second age of the RNA world; this second age lasted until the appearance of the first replicating DNA cell(s), which eliminated their parental RNA cell counterparts in Darwinian evolution (47). Even the catalytic component of present-day ribosomes is RNA, with proteins just providing a scaffold: “The ribosome is a ribozyme,” as has been vividly summarized (13). Thus, viroids comply with the lack of mRNA activity expected for a replicating relic of the (first age of the) RNA world, wherein no ribosomes existed.

Sixth and most important, some viroids display catalytic activity via hammerhead ribozymes. When Diener formulated his hypothesis (1989), ribozymes (the key signature of the RNA world) were known to be prevalent among viroid-like satellite RNAs but restricted to just one (and peculiar) viroid (ASBVd). Subsequent discovery of three additional viroids with hammerhead ribozymes in both polarity strands, PLMVd (71), CChMVd (93), and *Eggplant latent viroid* (39)—and more recently of two other candidates (144, 146)—dispelled the doubts that ASBVd could be a viroid-like satellite RNA and consolidated the family *Avsunviroidae*. The small size and relatively low structural complexity of the hammerhead and hairpin ribozymes, much simpler than those mediating intron splicing, also support their involvement in self-replication of RNA. The hammerhead and hairpin ribozymes can catalyze two of the three steps of viroid replication: cleavage and ligation, but not polymerization (**Figure 3**). RNA-catalyzed RNA polymerization has not been reported in nature, but in vitro evolution studies have shown that RNA can catalyze this reaction (77, 87, 142), thus making viroid replication in a preprotein world feasible.

Do viroids have a monophyletic origin? The low overall sequence similarity among these RNAs poses a problem for inferring a reliable phylogeny (76). Nevertheless, when alignments are adjusted—considering the local similarities and the insertions/deletions and duplications/rearrangements described—and an appropriate estimator of genetic distances is used, phylogenetic reconstructions support a monophyletic origin. Furthermore, these reconstructions are consistent with the major groups proposed previously for viroids (and viroid-like satellite RNAs), as well as with their biological properties (36, 37). Yet, a polyphyletic viroid origin cannot be dismissed, particularly considering the unusual base composition of ASBVd. Furthermore, thermodynamic predictions of the secondary structure of viroid RNAs support an evolutionary trend during viroid radiation toward increased robustness (resistance to deleterious mutations) and reduced antagonistic epistasis (interactions between deleterious mutations) (116, 117).

Transition from the RNA World to a World with Proteins and DNA: Viroids

Because nearly all intermediate stages have declined with time, it is difficult to speculate how viroids evolved from the RNA world, more than 3,000 mya, into a world involving two additional macromolecules (proteins and DNA). Nevertheless, at least some hints can be inferred from a close examination of these small replicating RNAs. Focusing on the distinction between families *Pospiviroidae* (or nuclear viroids) and *Avsunviroidae* (or chloroplastic viroids), two reasons argue in favor of the latter preceding the former. According to the endosymbiont hypothesis, plastids derive from primitive cyanobacteria by symbiosis (89); these free-living cells, which emerged prior to eukaryotic cells, might have hosted (and perhaps still host) the ancestors of the family *Avsunviroidae* before they adapted to plants (14, 86). In support of this view, the mediation of one step of the replication cycle by the hammerhead ribozyme has been retained only in the family *Avsunviroidae*. Conceivably, the most complex nuclear environment provided increased opportunities for the complete transition from RNA- to protein-catalyzed replication, whereas this transition has been partially accomplished in plastids. Yet, it is intriguing that the two families have adopted the same strategy regarding polymerization of RNA strands: to redirect DNA-dependent RNA polymerases to transcribe RNA templates. Although such a scenario may appear realistic for viroids replicating in plastids, wherein no RNA-dependent RNA polymerase has been described, the situation is different for viroids replicating in the nucleus, wherein several enzymes of this class accumulate. An alternative and more plausible explanation is that the DNA-like compact folding of the genomic viroid RNA (and its replicative intermediates) may have facilitated their recognition by RNA polymerases that transcribe double-stranded templates into ssRNAs, in contrast to RNA-dependent RNA polymerases that recognize ssRNA templates and convert them into dsRNA products.

On the one hand, two DNA-dependent RNA polymerases reside in plastids (1, 125), and the available evidence supports the involvement of one of them, the nuclear-encoded polymerase (NEP), in replication of members of the family *Avsunviroidae* (97, 113). This finding is consistent with the NEP, comprising a single subunit, being structurally simpler (and presumably older) than the plastid-encoded polymerase (PEP), comprising five subunits. On the other hand, the three nuclear DNA-dependent RNA polymerases typical of eukaryotes (RNA polymerases I to III) (17), and the two additional ones described recently in plants (RNA polymerases IV and V) (62), display a high level of structural complexity (with more than 10 subunits). Why members of the family *Pospiviroidae* have specifically co-opted RNA polymerase II (44, 92, 119) remains an enigma, but interestingly, the same enzyme is also involved in HDV RNA replication (129). Perhaps this aptitude of RNA polymerase II is related to its versatility in catalyzing transcription of protein-coding genes as well as long and small noncoding RNAs (62).

The RNA polymerases mediating replication of viroids are also the driving force that largely determines two of their features: the differential complexity of viroid quasi-species and the recombinant nature of some viroids. The mutation rate estimated for CChMVd, 0.0025 per site and replication cycle (one mutation per replicated genome), is the highest reported for any biological entity (**Figure 4**) (51). This extremely error-prone replication—most likely caused by a proofreading-deficient RNA polymerase (NEP) forced to use RNA instead of its native DNA template (43) and perhaps also by accumulation of mutagenic free radicals in plastids or unbalanced nucleotide pools—is reminiscent of that postulated for the primitive replicons of the RNA world. Viroids, and more explicitly members of the family *Avsunviroidae*, can possibly withstand this elevated mutation rate because they have a very small genome, whereas more complex genomes would accumulate an excessive mutational load (34). These results also suggest that emergence of mechanisms enhancing replication fidelity was critical for the evolution of complexity in the early




Figure 4

Per-site mutation rate versus genome size for *Chrysanthemum chlorotic mottle viroid* and other biological entities. RNA viruses (left to right) are *Tobacco mosaic virus*, human rhinovirus, poliovirus, *Vesicular stomatitis virus*, bacteriophage $\phi 6$, and *Measles virus*; all are single stranded except bacteriophage $\phi 6$, which is double stranded. Single-stranded DNA (ssDNA) viruses are bacteriophage $\phi X174$ and bacteriophage m13. Double-stranded DNA (dsDNA) viruses are bacteriophage λ , *Herpes simplex virus*, bacteriophage T2, and bacteriophage T4. Bacteria are *Escherichia coli*. Lower eukaryotes are *Saccharomyces cerevisiae* and *Neurospora crassa*. Higher eukaryotes are *Caenorhabditis elegans*, *Drosophila melanogaster*, *Mus musculus*, and *Homo sapiens*. Where several estimations were available, the mean value is shown. Reproduced with permission from Reference 51.

history of life (51). Furthermore, the need to transcribe atypical viroid RNA templates might also decrease the processivity of the otherwise DNA-dependent RNA polymerases, promoting their stalling and subsequent jumping with the bound nascent transcript, thus facilitating the appearance of chimeric viroids. Their recurrent identification (133) supports the view that recombination has played an important role in shaping the viroidsphere.

The other two replication steps further attest to the resilience of viroids. For cleavage of their oligomeric replication intermediates, members of the family *Pospiviroidae* recruit RNases of class III mediating posttranscriptional maturation of highly structured nuclear transcripts (53). Members of the *Avsunviroidae* recruit RNA-binding proteins with a similar role in processing of chloroplastic transcripts, which additionally behave as RNA chaperones that in vitro, and presumably in vivo, stimulate the efficiency of hammerhead ribozymes (20). For circularization of the resulting unit-length strands, PSTVd (and most likely the other members of the family *Pospiviroidae*) diverts DNA ligase 1 to function as an RNA ligase, taking advantage of the fact that, upon DNA binding, DNA ligase 1 promotes a local RNA-like A-form conformation in the vicinity of the DNA break (102, 103). Moreover, DNA ligase 1 and some RNA ligases, like T4 RNA ligase 1, demand 5'-phosphomonoester and 3'-hydroxyl termini like those generated by class III RNases and operate through a similar mechanism (72, 137). Further supporting this view, DNA ligases, RNA ligases, and RNA-capping enzymes probably evolved by fusion between auxiliary effector domains and an ancestral catalytic module mediating RNA repair (123). Thus, some viroids would opportunistically exploit the preservation of a primordial role of DNA ligase 1 (98). In contrast, for ligation of the (+) and (-) monomeric linear replication intermediates, members of the family *Avsunviroidae* recruit a chloroplastic tRNA ligase isoform with specificity for the 5'-hydroxyl and 2',3' cyclic phosphodiester termini produced by hammerhead ribozymes (99).

Transition from the RNA World to a World with Proteins and DNA: Viroid-Related Replicons

The features of viroid-like satellite RNAs, both structural (small size and circularity) and functional (lack of mRNA ability and replication by a rolling-circle mechanism mediated by hammerhead and hairpin ribozymes), support a common origin with viroids and, particularly, with members of the family *Avsunviroidae* (36). However, in their subsequent evolution, viroid-like satellite RNAs have followed a different strategy: Instead of relying exclusively on the host transcriptional and processing machinery, they have additionally parasitized a helper virus that provides the transmission and most likely part of the replication machinery. This ménage à trois, more complex but possibly more efficient for ensuring dispersal of the viroid-like RNAs by the helper virus—viroids are self-sufficient in this respect—also suggests that parasitism emerged very early in the history of life.

Resembling viroid-like satellite RNAs, HDV has also evolved to parasitize a helper virus, but only for transmission because HDV RNA, like viroids, replicates autonomously in infected cells. How HDV RNA might have appeared has been discussed in extenso previously (130). Here we just emphasize that analysis of HDV RNA has uncovered its recombinant nature: The rodlike secondary structure can be regarded as the fusion of a viroid-like domain, holding the ribozymes and located in one of the terminal domains, to another domain that in the antigenomic polarity contains the open reading frame for the so-called δ antigen (δ Ag), a protein with a crucial role in the biology of the virus (60). This modular design has led to the proposal that HDV RNA might have arisen from a self-replicating, viroid-like RNA capturing the mRNA encoding a protein, like the δ Ag, which, having a beneficial effect on the recombinant product, would have secured its persistence (8, 112).

In contrast to viroids and the other viroid-related replicons, retroviroid-like elements have a DNA counterpart and, therefore, should have appeared following the “invention” of this biopolymer. DNA can be viewed as a derivative of RNA, since all organisms synthesize deoxyribonucleotides by reducing ribonucleotides, and thymine by methylating uracil (85). The first of these two reactions, due to its complex sulfur-based mechanism, seems unlikely to have been catalyzed by a ribozyme, thus demanding a protein enzyme. This argument and others (49, 85) support the view that DNA appeared during the second age of the RNA world (after the advent of the ribosome) (47). RT, mediating transition from RNA to DNA, most likely evolved from an ancestral RNA-dependent RNA polymerase—ribozyme or protein enzyme—involved in replication of early RNA genomes (91, 104, 145). By acquiring a homologous DNA from which RNA can be transcribed, retroviroid-like elements would have increased their chances for survival. Intriguingly, following initial discovery of hammerhead ribozymes in certain newt transcripts (38), recent studies have unveiled the existence of hammerhead and HDV-like ribozymes along the biological scale, with the insertion site of the corresponding DNAs in the host genomes indicating a possible regulatory function and the contribution of retrotransposition to their genesis (22, 65, 140). However, it is unclear whether the RNA products resulting from ribozyme-mediated self-cleavage are subsequently circularized.

SUMMARY

The time had clearly come to ask how the DNA → RNA → protein flow of information had ever gotten started. Here, Francis [Crick] was again far ahead of his time. In 1968 he argued that RNA must have been the first genetic molecule, further suggesting that RNA, besides acting as a template, might also act as an enzyme and, in so doing, catalyze its own self-replication.

—J.D. Watson (54, p. xxiii)

The course of time has provided accumulating evidence to this passionate laudation by J.D. Watson in the prologue of the first edition (54) of the classic book on the RNA world (3). Just a recent example: The synthesis of activated pyrimidine ribonucleotides starting from plausible prebiotic precursors and conditions has bypassed the difficulty of how ribonucleotides could have been formed from their constituent parts (ribose and nucleobases) and removed another important hurdle for the emergence of the RNA world (105). The survival until today of viroids (and viroid-related replicons) adds further support to this view.

1. Viroids fulfill all structural criteria postulated for primitive replicons, including small size, (G + C)-rich content and compact folding, as well as circularity that may ensure complete replication and even polyploidy.
2. From a functional perspective, viroids lack protein-coding ability, but at least some encode ribozymes; in other words, they are catalytic RNAs, the key signature expected for survivors of the RNA world.
3. The three catalytic activities mediating rolling-circle replication of viroids, namely, RNA polymerization, cleavage, and ligation, can be supplied by natural or artificial ribozymes, making replication reasonable in a protein-free world.
4. Viroid-like satellite RNAs, retroviroid-like elements, and HDV RNA can be regarded as evolutionary inventions related to (and possibly derived from) viroids.
5. Feasible evolutionary routes can be envisaged for the transition of viroids and viroid-related replicons from the RNA world to a world with proteins and DNA.

Science runs through winding roads. The initial purpose of characterizing the viruses presumed to cause several plant diseases eventually, and unexpectedly, resulted in the discovery of a new class of subviral agents, the viroids. Their properties are consistent with those predicted for the first minimal replicons that populated the RNA world, bolstering the case for its existence.

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

R.F. has received funding by grant BFU2011-28443 from Ministerio de Economía y Competitividad (MINECO, Spain), R.S. by grants BFU2011-25271 (MINECO) and ERC-2011-StG-281191-VIRMUT (European Research Council), and S.F.E. by grant BFU2012-30805 (MINECO). P.S. has been supported by postdoctoral contracts from Generalitat Valenciana (APOSTD/2010, program VALi+d) and MINECO (program Juan de la Cierva).

LITERATURE CITED

1. Allison LA, Simon LD, Maliga P. 1996. Deletion of *rpoB* reveals a second distinct transcription system in plastids of higher plants. *EMBO J.* 15:2802–9
2. Ambrós S, Hernández C, Flores R. 1999. Rapid generation of genetic heterogeneity in progenies from individual cDNA clones of peach latent mosaic viroid in its natural host. *J. Gen. Virol.* 80:2239–52
3. Atkins JF, Gesteland RF, Cech TR, eds. 2011. *RNA Worlds: From Life's Origins to Diversity in Gene Regulation*. Cold Spring Harbor, NY: Cold Spring Harbor Lab. Press
4. Beijerinck MW. 1898. Ueber ein Contagium vivum fluidum als Ursache der Fleckenkrankheit der Tabaksblätter. *Verh. K. Akad. Wet. Amst.* 65:3–21

5. Bernad L, Duran-Vila N, Elena SF. 2009. Effect of citrus hosts on the generation, maintenance and evolutionary fate of genetic variability of citrus exocortis viroid. *J. Gen. Virol.* 90:2040–9
6. Biebricher CK, Eigen M. 2006. What is a quasispecies? *Curr. Top. Microbiol. Immunol.* 299:1–31
7. Branch AD, Robertson HD. 1984. A replication cycle for viroids and other small infectious RNAs. *Science* 223:450–4
8. Brazas R, Ganem D. 1996. A cellular homolog of hepatitis delta antigen: implications for viral replication and evolution. *Science* 274:90–4
9. Briones C, Stich M, Manrubia SC. 2009. The dawn of the RNA world: toward functional complexity through ligation of random RNA oligomers. *RNA* 15:743–49
10. Bruening G, Passmore BK, van Tol H, Buzayan JM, Feldstein PA. 1991. Replication of a plant virus satellite RNA: Evidence favors transcription of circular templates of both polarities. *Mol. Plant Microbe Interact.* 4:219–25
11. Burch CL, Chao L. 1999. Evolution by small steps and rugged landscapes in the RNA virus $\phi 6$. *Genetics* 151:921–27
12. Buzayan JM, Gerlach WL, Bruening G. 1986. Non-enzymatic cleavage and ligation of RNAs complementary to a plant virus satellite RNA. *Nature* 323:349–53
13. Cech TR. 2000. The ribosome is a ribozyme. *Science* 289:878–79
14. Chela-Flores J. 1994. Are viroids molecular fossils of the RNA world? *J. Theor. Biol.* 166:163–66
15. Chen PJ, Kalpana G, Goldberg J, Mason W, Werner B, et al. 1986. Structure and replication of the genome of the hepatitis δ virus. *Proc. Natl. Acad. Sci. USA* 83:8774–78
16. Codoñer FM, Daròs JA, Solé RV, Elena SF. 2006. The fittest versus the flattest: experimental confirmation of the quasispecies effect with subviral pathogens. *PLoS Pathog.* 2:1187–93
17. Cramer P, Armache KJ, Baumli S, Benkert S, Brueckner F, et al. 2008. Structure of eukaryotic RNA polymerases. *Annu. Rev. Biophys.* 37:337–52
18. Crick FH. 1968. The origin of the genetic code. *J. Mol. Biol.* 38:367–79
19. Daròs JA, Flores R. 1995. Identification of a retroviroid-like element from plants. *Proc. Natl. Acad. Sci. USA* 92:6856–60
20. Daròs JA, Flores R. 2002. A chloroplast protein binds a viroid RNA in vivo and facilitates its hammerhead-mediated self-cleavage. *EMBO J.* 21:749–59
21. Daròs JA, Marcos JF, Hernández C, Flores R. 1994. Replication of avocado sunblotch viroid: evidence for a symmetric pathway with two rolling circles and hammerhead ribozyme processing. *Proc. Natl. Acad. Sci. USA* 91:12813–17
22. de la Peña M, García-Robles I. 2010. Ubiquitous presence of the hammerhead ribozyme motif along the tree of life. *RNA* 16:1943–50
23. de la Peña M, Navarro B, Flores R. 1999. Mapping the molecular determinant of pathogenicity in a hammerhead viroid: a tetraloop within the in vivo branched RNA conformation. *Proc. Natl. Acad. Sci. USA* 96:9960–65
24. Delgado S, Martínez de Alba AE, Hernández C, Flores R. 2005. A short double-stranded RNA motif of peach latent mosaic viroid contains the initiation and the self-cleavage sites of both polarity strands. *J. Virol.* 79:12934–43
25. den Boon JA, Ahlquist P. 2010. Organelle-like membrane compartmentalization of positive-strand RNA virus replication factories. *Annu. Rev. Microbiol.* 64:241–56
26. Diener TO. 1971. Potato spindle tuber “virus”: IV. A replicating, low molecular weight RNA. *Virology* 45:411–28
27. Diener TO. 1972. Potato spindle tuber viroid: VIII. Correlation of infectivity with a UV-absorbing component and thermal denaturation properties of the RNA. *Virology* 50:606–9
28. Diener TO. 1981. Are viroids escaped introns? *Proc. Natl. Acad. Sci. USA* 78:5014–15
29. Diener TO. 1989. Circular RNAs: relics of precellular evolution? *Proc. Natl. Acad. Sci. USA* 86:9370–74
30. Diener TO. 2003. Discovering viroids: a personal perspective. *Nat. Rev. Microbiol.* 1:75–80
31. Ding B. 2009. The biology of viroid-host interactions. *Annu. Rev. Phytopathol.* 47:105–31
32. Ding B. 2010. Viroids: self-replicating, mobile, and fast-evolving noncoding regulatory RNAs. *Wiley Interdiscip. Rev. RNA* 1:362–75

33. Domingo E, Sabo D, Taniguchi T, Weissmann C. 1978. Nucleotide sequence heterogeneity of an RNA phage population. *Cell* 13:735–44
34. Eigen M. 1971. Self-organization of matter and the evolution of biological macromolecules. *Naturwissenschaften* 58:465–523
35. Eigen M, Schuster P. 1978. The hypercycle. A principle of natural self-organization. Part C: the realistic hypercycle. *Naturwissenschaften* 65:341–69
36. Elena SF, Dopazo J, de la Peña M, Flores R, Diener TO, Moya A. 2001. Phylogenetic analysis of viroid and viroid-like satellite RNAs from plants: a reassessment. *J. Mol. Evol.* 53:155–59
37. Elena SF, Dopazo J, Flores R, Diener TO, Moya A. 1991. Phylogeny of viroids, viroidlike satellite RNAs, and the viroidlike domain of hepatitis δ virus RNA. *Proc. Natl. Acad. Sci. USA* 88:5631–34
38. Epstein LM, Gall JG. 1987. Self-cleaving transcripts of satellite DNA from the newt. *Cell* 48:535–43
39. Fadda Z, Daròs JA, Fagoaga C, Flores R, Duran-Vila N. 2003. *Eggplant latent viroid*, the candidate type species for a new genus within the family *Ausunviroidae* (hammerhead viroids). *J. Virol.* 77:6528–32
40. Fedor MJ. 2000. Structure and function of the hairpin ribozyme. *J. Mol. Biol.* 297:269–91
41. Feix G, Pollet R, Weissmann C. 1968. Replication of viral RNA, XVI. Enzymatic synthesis of infectious viral RNA with noninfectious Q-beta minus strands as template. *Proc. Natl. Acad. Sci. USA* 59:145–52
42. Flores R, Di Serio F, Navarro B, Duran-Vila N, Owens RA. 2011. Viroids and viroid diseases of plants. In *Studies in Viral Ecology, Vol. I: Microbial and Botanical Host Systems*, ed. CJ Hurst, pp. 311–46. Hoboken, NJ: John Wiley & Sons
43. Flores R, Hernández C, Martínez de Alba E, Daròs JA, Di Serio F. 2005. Viroids and viroid-host interactions. *Annu. Rev. Phytopathol.* 43:117–39
44. Flores R, Semancik JS. 1982. Properties of a cell-free system for synthesis of citrus exocortis viroid. *Proc. Natl. Acad. Sci. USA* 79:6285–88
45. Flores R, Serra P, Minoia S, Di Serio F, Navarro B. 2012. Viroids: from genotype to phenotype just relying on RNA sequence and structural motifs. *Front. Microbiol.* 3:217
46. Forster AC, Symons RH. 1987. Self-cleavage of plus and minus RNAs of a virusoid and a structural model for the active sites. *Cell* 49:211–20
47. Forterre P. 2005. The two ages of the RNA world, and the transition to the DNA world: a story of viruses and cells. *Biochimie* 87:793–803
48. Fraenkel-Conrat H. 1956. The role of the nucleic acid in the reconstitution of active tobacco mosaic virus. *J. Am. Chem. Soc.* 78:882–83
49. Freeland SJ, Knight RD, Landweber LF. 1999. Do proteins predate DNA? *Science* 286:690–92
50. Fresco JR, Alberts BM, Doty P. 1960. Some molecular details of the secondary structure of ribonucleic acid. *Nature* 188:98–101
51. Gago S, Elena SF, Flores R, Sanjuán R. 2009. Extremely high mutation rate of a hammerhead viroid. *Science* 323:1308
52. Gandía M, Duran-Vila N. 2004. Variability of the progeny of a sequence variant *Citrus bent leaf viroid* (CBLVd). *Arch. Virol.* 149:407–16
53. Gas ME, Hernández C, Flores R, Daròs JA. 2007. Processing of nuclear viroids in vivo: an interplay between RNA conformations. *PLoS Pathog.* 3:1813–26
54. Gesteland RF, Atkins JF, eds. 1993. *The RNA World*. Cold Spring Harbor, NY: Cold Spring Harbor Lab. Press
55. Gierer A, Schramm G. 1956. Infectivity of ribonucleic acid from tobacco mosaic virus. *Nature* 177:702–3
56. Gilbert W. 1986. Origin of life: the RNA world. *Nature* 319:618
57. Góra-Sochacka A, Kierzek A, Candresse T, Zagórski W. 1997. The genetic stability of potato spindle tuber viroid (PSTVd) molecular variants. *RNA* 3:68–74
58. Grill LK, Semancik JS. 1978. RNA sequences complementary to citrus exocortis viroid in nucleic acid preparations from infected *Gynura aurantiaca*. *Proc. Natl. Acad. Sci. USA* 75:896–900
59. Gross HJ, Domdey H, Lossow C, Jank P, Raba M, et al. 1978. Nucleotide sequence and secondary structure of potato spindle tuber viroid. *Nature* 273:203–8
60. Gudima S, Wu SY, Chiang CM, Moraleda G, Taylor J. 2000. Origin of hepatitis δ virus mRNA. *J. Virol.* 74:7204–10

61. Guerrier-Takada C, Gardiner K, Marsh T, Pace N, Altman S. 1983. The RNA moiety of ribonuclease P is the catalytic subunit of the enzyme. *Cell* 35:849–57
62. Haag JR, Pikaard CS. 2011. Multisubunit RNA polymerases IV and V: purveyors of non-coding RNA for plant gene silencing. *Nat. Rev. Mol. Cell Biol.* 12:483–92
63. Hadidi A. 1986. Relationship of viroids and certain other plant pathogenic nucleic acids to group I and II introns. *Plant Mol. Biol.* 7:129–42
64. Hajeri S, Ramadugu C, Manjunath K, Ng J, Lee R, Vidalakis G. 2011. In vivo generated *Citrus exocortis* viroid progeny variants display a range of phenotypes with altered levels of replication, systemic accumulation and pathogenicity. *Virology* 417:400–9
65. Hammann C, Luptak A, Perreault J, de la Peña M. 2012. The ubiquitous hammerhead ribozyme. *RNA* 18:871–85
66. Hammann C, Steger G. 2012. Viroid-specific small RNA in plant disease. *RNA Biol.* 9:809–19
67. Hammond R, Smith DR, Diener TO. 1989. Nucleotide sequence and proposed secondary structure of *Columnea* latent viroid: a natural mosaic of viroid sequences. *Nucleic Acids Res.* 17:10083–94
68. Haruna I, Spiegelman S. 1966. A search for an intermediate involving a complement during synchronous synthesis by a purified RNA replicase. *Proc. Natl. Acad. Sci. USA* 55:1256–63
69. Hegedus K, Dallmann G, Balazs E. 2004. The DNA form of a retroviroid-like element is involved in recombination events with itself and with the plant genome. *Virology* 325:277–86
70. Hernández C, Daròs JA, Elena SF, Moya A, Flores R. 1992. The strands of both polarities of a small circular RNA from carnation self-cleave in vitro through alternative double- and single-hammerhead structures. *Nucleic Acids Res.* 20:6323–29
71. Hernández C, Flores R. 1992. Plus and minus RNAs of peach latent mosaic viroid self-cleave in vitro via hammerhead structures. *Proc. Natl. Acad. Sci. USA* 89:3711–15
72. Ho CK, Shuman S. 2002. Bacteriophage T4 RNA ligase 2 (gp24.1) exemplifies a family of RNA ligases found in all phylogenetic domains. *Proc. Natl. Acad. Sci. USA* 99:12709–14
73. Hutchins CJ, Keese P, Visvader JE, Rathjen PD, McInnes JL, Symons RH. 1985. Comparison of multimeric plus and minus forms of viroids and virusoids. *Plant Mol. Biol.* 4:293–304
74. Hutchins CJ, Rathjen PD, Forster AC, Symons RH. 1986. Self-cleavage of plus and minus RNA transcripts of avocado sunblotch viroid. *Nucleic Acids Res.* 14:3627–40
75. Ivica NA, Obermayer B, Campbell GW, Rajamani S, Gerland U, Chen IA. 2013. The paradox of dual roles in the RNA world: resolving the conflict between stable folding and templating ability. *J. Mol. Evol.* 77:55–63
76. Jenkins GM, Woelk CH, Rambaut A, Holmes EC. 2000. Testing the extent of sequence similarity among viroids, satellite RNAs, and the hepatitis delta virus. *J. Mol. Evol.* 50:98–102
77. Johnston WK, Unrau PJ, Lawrence MS, Glasner ME, Bartel DP. 2001. RNA-catalyzed RNA polymerization: accurate and general RNA-templated primer extension. *Science* 292:1319–25
78. Juhasz A, Hegyi H, Solymosy F. 1988. A novel aspect of the information content of viroids. *Biochim. Biophys. Acta* 950:455–58
79. Kiefer MC, Owens RA, Diener TO. 1983. Structural similarities between viroids and transposable genetic elements. *Proc. Natl. Acad. Sci. USA* 80:6234–38
80. Kolonko N, Bannach O, Aschermann K, Hu KH, Moors M, et al. 2006. Transcription of potato spindle tuber viroid by RNA polymerase II starts in the left terminal loop. *Virology* 347:392–404
81. Kruger K, Grabowski PJ, Zaugg AJ, Sands J, Gottschling DE, Cech TR. 1982. Self-splicing RNA: auto-excision and autocyclization of the ribosomal RNA intervening sequence of *Tetrahymena*. *Cell* 31:147–57
82. Kuo MY, Goldberg J, Coates L, Mason W, Gerin J, Taylor J. 1988. Molecular cloning of hepatitis delta virus RNA from an infected woodchuck liver: sequence, structure, and applications. *J. Virol.* 62:1855–61
83. Kuo MY, Sharmeen L, Dinter-Gottlieb G, Taylor J. 1988. Characterization of self-cleaving RNA sequences on the genome and antigenome of human hepatitis delta virus. *J. Virol.* 62:4439–44
84. Lauring AS, Andino R. 2010. Quasispecies theory and the behavior of RNA viruses. *PLoS Pathog.* 6:e1001005
85. Lazcano A, Guerrero R, Margulis L, Oro J. 1988. The evolutionary transition from RNA to DNA in early cells. *J. Mol. Evol.* 27:283–90

86. Lima MI, Fonseca MEN, Flores R, Kitajima EW. 1994. Detection of avocado sunblotch viroid in chloroplasts of avocado leaves by in situ hybridization. *Arch. Virol.* 138:385–90
87. Lincoln TA, Joyce GF. 2009. Self-sustained replication of an RNA enzyme. *Science* 323:1229–32
88. Manrubia SC, Briones C. 2007. Modular evolution and increase of functional complexity in replicating RNA molecules. *RNA* 13:97–107
89. Margulis L. 1993. *Symbiosis in Cell Evolution*. New York: W.H. Freeman. 2nd ed.
90. McInnes JL, Symons RH. 1991. Comparative structure of viroids and their rapid detection using radioactive and nonradioactive nucleic acid probes. In *Viroids and Satellites: Molecular Parasites at the Frontier of Life*, ed. K Maramorosch, pp. 21–58. Boca Raton, FL: CRC
91. Moelling K. 2013. What contemporary viruses tell us about evolution: a personal view. *Arch. Virol.* 158:1833–48
92. Mühlbach HP, Sängler HL. 1979. Viroid replication is inhibited by α -amanitin. *Nature* 278:185–88
93. Navarro B, Flores R. 1997. Chrysanthemum chlorotic mottle viroid: unusual structural properties of a subgroup of viroids with hammerhead ribozymes. *Proc. Natl. Acad. Sci. USA* 94:11262–67
94. Navarro B, Gisel A, Rodio ME, Delgado S, Flores R, Di Serio F. 2012. Small RNAs containing the pathogenic determinant of a chloroplast-replicating viroid guide the degradation of a host mRNA as predicted by RNA silencing. *Plant J.* 70:991–1003
95. Navarro B, Gisel A, Rodio ME, Delgado S, Flores R, Di Serio F. 2012. Viroids: how to infect a host and cause disease without encoding proteins. *Biochimie* 94:1474–80
96. Navarro JA, Flores R. 2000. Characterization of the initiation sites of both polarity strands of a viroid RNA reveals a motif conserved in sequence and structure. *EMBO J.* 19:2662–70
97. Navarro JA, Vera A, Flores R. 2000. A chloroplastic RNA polymerase resistant to tagetitoxin is involved in replication of *Avocado sunblotch viroid*. *Virology* 268:218–25
98. Nohales MA, Flores R, Daròs JA. 2012. A viroid RNA redirects host DNA ligase 1 to act as an RNA ligase. *Proc. Natl. Acad. Sci. USA* 109:13805–10
99. Nohales MA, Molina-Serrano D, Flores R, Daròs JA. 2012. Involvement of the chloroplastic isoform of tRNA ligase in the replication of the viroids belonging to the family *Avsunviroidae*. *J. Virol.* 86:8269–76
100. Ojosnegros S, Perales C, Mas A, Domingo E. 2011. Quasispecies as a matter of fact: viruses and beyond. *Virus Res.* 162:203–15
101. Orgel LE. 1968. Evolution of the genetic apparatus. *J. Mol. Biol.* 38:381–93
102. Pascal JM. 2008. DNA and RNA ligases: Structural variations and shared mechanisms. *Curr. Opin. Struct. Biol.* 18:96–105
103. Pascal JM, O'Brien PJ, Tomkinson AE, Ellenberger T. 2004. Human DNA ligase I completely encircles and partially unwinds nicked DNA. *Nature* 432:473–78
104. Poole AM, Logan DT. 2005. Modern mRNA proofreading and repair: clues that the last universal common ancestor possessed an RNA genome? *Mol. Biol. Evol.* 22:1444–55
105. Powner MW, Gerland B, Sutherland JD. 2009. Synthesis of activated pyrimidine ribonucleotides in prebiotically plausible conditions. *Nature* 459:239–42
106. Prody GA, Bakos JT, Buzayan JM, Schneider IR, Bruening G. 1986. Autolytic processing of dimeric plant virus satellite RNA. *Science* 231:1577–80
107. Reaney D. 1984. Genetic noise in evolution? *Nature* 307:318–19
108. Rezaian MA. 1990. Australian grapevine viroid—evidence for extensive recombination between viroids. *Nucleic Acids Res.* 18:1813–18
109. Riesner D, Henco K, Rokohl U, Klotz G, Kleinschmidt AK, et al. 1979. Structure and structure formation of viroids. *J. Mol. Biol.* 133:85–115
110. Rizzetto M, Canese MG, Aricò J, Crivelli O, Trepo C, et al. 1977. Immunofluorescence detection of a new antigen-antibody system (δ /anti- δ) associated to the hepatitis B virus in the liver and in the serum of HBsAg carriers. *Gut* 18:997–1003
111. Rizzetto M, Hoyer B, Canese MG, Shih JW, Purcell RH, Gerin JL. 1980. δ agent: association of δ antigen with hepatitis B surface antigen and RNA in serum of δ -infected chimpanzees. *Proc. Natl. Acad. Sci. USA* 77:6124–28
112. Robertson HD. 1996. How did replicating and coding RNAs first get together? *Science* 274:66–67

113. Rodio ME, Delgado S, de Stradis AE, Gómez MD, Flores R, Di Serio F. 2007. A viroid RNA with a specific structural motif inhibits chloroplast development. *Plant Cell* 19:3610–26
114. Roossinck MJ, Sleat D, Palukaitis P. 1992. Satellite RNAs of plant viruses: structures and biological effects. *Microbiol. Rev.* 56:265–79
115. Sanjuán R, Cuevas JM, Furió V, Holmes EC, Moya A. 2007. Selection for robustness in mutagenized RNA viruses. *PLoS Genet.* 3:939–46
116. Sanjuán R, Forment J, Elena SF. 2006. In silico predicted robustness of viroid RNA secondary structures. II. Interaction between mutation pairs. *Mol. Biol. Evol.* 23:2123–30
117. Sanjuán R, Forment J, Elena SF. 2006. In silico predicted robustness of viroids RNA secondary structures. I. The effect of single mutations. *Mol. Biol. Evol.* 23:1427–36
118. Schiebel W, Pélissier T, Riedel L, Thalmeir S, Schiebel R, et al. 1998. Isolation of an RNA-directed RNA polymerase-specific cDNA clone from tomato. *Plant Cell* 10:2087–101
119. Schindler IM, Mühlbach HP. 1992. Involvement of nuclear DNA-dependent RNA polymerases in potato spindle tuber viroid replication: a reevaluation. *Plant Sci.* 84:221–29
120. Schneider R. 1969. Satellite-like particle of tobacco ringspot virus that resembles tobacco ringspot virus. *Science* 166:1627–29
121. Schuster P, Swetina J. 1988. Stationary mutant distributions and evolutionary optimization. *Bull. Math. Biol.* 50:635–60
122. Semancik JS, Szychowski JA, Rakowski AG, Symons RH. 1993. Isolates of citrus exocortis viroid recovered by host and tissue selection. *J. Gen. Virol.* 74:2427–36
123. Shuman S, Lima CD. 2004. The polynucleotide ligase and RNA capping enzyme superfamily of covalent nucleotidyltransferases. *Curr. Opin. Struct. Biol.* 14:757–64
124. Sogo JM, Koller T, Diener TO. 1973. Potato spindle tuber viroid. X: Visualization and size determination by electron microscopy. *Virology* 55:70–80
125. Sugiura M. 1992. The chloroplast genome. *Plant Mol. Biol.* 19:149–68
126. Symons RH. 1981. Avocado sunblotch viroid: primary sequence and proposed secondary structure. *Nucleic Acids Res.* 9:6527–37
127. Symons RH, Randles JW. 1999. Encapsidated circular viroidlike satellite RNAs (virusoids) of plants. *Curr. Top. Microbiol. Immunol.* 239:81–105
128. Tabler M, Tsagris M. 2004. Viroids: petite RNA pathogens with distinguished talents. *Trends Plant Sci.* 9:339–48
129. Taylor JM. 2009. Replication of the hepatitis delta virus RNA genome. *Adv. Virus Res.* 74:103–21
130. Taylor JM, Pelchat M. 2010. Origin of hepatitis δ virus. *Future Microbiol.* 5:393–402
131. Tessitori M, Rizza S, Reina A, Causarano G, Di Serio F. 2013. The genetic diversity of *Citrus dwarfing viroid* populations is mainly dependent on the infected host species. *J. Gen. Virol.* 94:687–93
132. Tsagris EM, Martínez de Alba AE, Gozmanova M, Kalantidis K. 2008. Viroids. *Cell Microbiol.* 10:2168–79
133. Verhoeven JT, Meeke ET, Roenhorst JW, Flores R, Serra P. 2013. Dahlia latent viroid: a recombinant new species of the family *Pospiviroidae* posing intriguing questions about its origin and classification. *J. Gen. Virol.* 94:711–19
134. Vignuzzi M, Stone JK, Arnold JJ, Cameron CE, Andino R. 2006. Quasispecies diversity determines pathogenesis through cooperative interactions in a viral population. *Nature* 439:344–48
135. Visvader JE, Symons RH. 1985. Eleven new sequence variants of citrus exocortis viroid and the correlation of sequence with pathogenicity. *Nucleic Acids Res.* 13:2907–20
136. Wang KS, Choo QL, Weiner AJ, Ou JH, Najarian RC, et al. 1986. Structure, sequence and expression of the hepatitis delta (δ) viral genome. *Nature* 323:508–14. Corrigendum. 1987. *Nature* 328:456
137. Wang LK, Schwer B, Shuman S. 2006. Structure-guided mutational analysis of T4 RNA ligase 1. *RNA* 12:2126–34
138. Wassenegger M, Krczal G. 2006. Nomenclature and functions of RNA-directed RNA polymerases. *Trends Plant Sci.* 11:142–51
139. Wassenegger M, Spieker RL, Thalmeir S, Gast FU, Riedel L, Sanger HL. 1996. A single nucleotide substitution converts potato spindle tuber viroid (PSTVd) from a noninfectious to an infectious RNA for *Nicotiana tabacum*. *Virology* 226:191–97

140. Webb C-HT, Riccitelli NJ, Ruminski DJ, Lupták A. 2009. Widespread occurrence of self-cleaving ribozymes. *Science* 326:953
141. Wilke CO, Wang JL, Ofria C, Lenski RE, Adami C. 2001. Evolution of digital organisms at high mutation rates leads to survival of the flattest. *Nature* 412:331–33
142. Wochner A, Attwater J, Coulson A, Holliger P. 2011. Ribozyme-catalyzed transcription of an active ribozyme. *Science* 332:209–12
143. Woese CR. 1967. *The Genetic Code: The Molecular Basis for Genetic Expression*. New York: Harper and Row
144. Wu Q, Wang Y, Cao M, Pantaleo V, Burgyan J, et al. 2012. Homology-independent discovery of replicating pathogenic circular RNAs by deep sequencing and a new computational algorithm. *Proc. Natl. Acad. Sci. USA* 109:3938–43
145. Xiong Y, Eickbush TH. 1990. Origin and evolution of retroelements based upon their reverse transcriptase sequences. *EMBO J.* 9:3353–62
146. Zhang ZX, Chen SS, Qi SS, Ding SW, Wu QF, Li SF. 2013. *Identification of new viroid-like circular RNAs from grapevine and apple plants*. Presented at Int. Workshop on Viroids and Satell. RNAs. Beijing
147. Zhong X, Archual AJ, Amin AA, Ding B. 2008. A genomic map of viroid RNA motifs critical for replication and systemic trafficking. *Plant Cell* 20:35–47



Contents

Beyond My Wildest Expectations <i>Eugene Nester</i>	1
Viral Miniproteins <i>Daniel DiMaio</i>	21
6S RNA, a Global Regulator of Transcription in <i>Escherichia coli</i> , <i>Bacillus subtilis</i> , and Beyond <i>Amy T. Cavanagh and Karen M. Wassarman</i>	45
Taming Wild Yeast: Potential of Conventional and Nonconventional Yeasts in Industrial Fermentations <i>Jan Steensels and Kevin J. Verstrepen</i>	61
Lipoteichoic Acid Synthesis and Function in Gram-Positive Bacteria <i>Matthew G. Percy and Angelika Gründling</i>	81
Temperature Sensing by Membranes <i>Diego de Mendoza</i>	101
What Ecologists Can Tell Virologists <i>John J. Dennehy</i>	117
The Medium Is the Message: Interspecies and Interkingdom Signaling by Peptidoglycan and Related Bacterial Glycans <i>Jonathan Dworkin</i>	137
Prokaryotic Ubiquitin-Like Protein Modification <i>Julie A. Maupin-Furlow</i>	155
The Importance of Microbes in Animal Development: Lessons from the Squid-Vibrio Symbiosis <i>Margaret J. McFall-Ngai</i>	177
The Tiniest Tiny Genomes <i>Nancy A. Moran and Gordon M. Bennett</i>	195
Effects of Antibiotics on Human Microbiota and Subsequent Disease <i>Kristie M. Keeney, Sophie Yurist-Doutsch, Marie-Claire Arrieta, and B. Brett Finlay</i>	217

Recombination Promoted by DNA Viruses: Phage λ to Herpes Simplex Virus <i>Sandra K. Weller and James A. Sawitzke</i>	237
The Peculiarities and Paradoxes of <i>Plasmodium</i> Heme Metabolism <i>Paul A. Sigala and Daniel E. Goldberg</i>	259
Biomass Utilization by Gut Microbiomes <i>Bryan A. White, Raphael Lamed, Edward A. Bayer, and Harry J. Flint</i>	279
Altered Egos: Antibiotic Effects on Food Animal Microbiomes <i>Heather K. Allen and Thad B. Stanton</i>	297
<i>Salmonella enterica</i> Serovar Typhi and the Pathogenesis of Typhoid Fever <i>Gordon Dougan and Stephen Baker</i>	317
Friend Turned Foe: Evolution of Enterococcal Virulence and Antibiotic Resistance <i>Daria Van Tyne and Michael S. Gilmore</i>	337
Bacterial Sigma Factors: A Historical, Structural, and Genomic Perspective <i>Andrey Feklistov, Brian D. Sharon, Seth A. Darst, and Carol A. Gross</i>	357
Fungal Membrane Organization: The Eisosome Concept <i>Lois M. Douglas and James B. Konopka</i>	377
Viroids: Survivors from the RNA World? <i>Ricardo Flores, Selma Gago-Zachert, Pedro Serra, Rafael Sanjuán, and Santiago F. Elena</i>	395
Bacterial Type III Secretion Systems: Specialized Nanomachines for Protein Delivery into Target Cells <i>Jorge E. Galán, Maria Lara-Tejero, Thomas C. Marlovits, and Samuel Wagner</i>	415
Subterfuge and Sabotage: Evasion of Host Innate Defenses by Invasive Gram-Positive Bacterial Pathogens <i>Cheryl Y.M. Okumura and Victor Nizet</i>	439
Fluorescence Imaging for Bacterial Cell Biology: From Localization to Dynamics, From Ensembles to Single Molecules <i>Zhibzhong Yao and Rut Carballido-López</i>	459
Cellular Sensing of Viral DNA and Viral Evasion Mechanisms <i>Megan H. Orzalli and David M. Knipe</i>	477
Regulation of Virulence of <i>Entamoeba histolytica</i> <i>Chelsea Marie and William A. Petri Jr.</i>	493



ANNUAL REVIEWS

It's about time. Your time. It's time well spent.

New From Annual Reviews:

Annual Review of Statistics and Its Application

Volume 1 • Online January 2014 • <http://statistics.annualreviews.org>

Editor: **Stephen E. Fienberg**, *Carnegie Mellon University*

Associate Editors: **Nancy Reid**, *University of Toronto*

Stephen M. Stigler, *University of Chicago*

The *Annual Review of Statistics and Its Application* aims to inform statisticians and quantitative methodologists, as well as all scientists and users of statistics about major methodological advances and the computational tools that allow for their implementation. It will include developments in the field of statistics, including theoretical statistical underpinnings of new methodology, as well as developments in specific application domains such as biostatistics and bioinformatics, economics, machine learning, psychology, sociology, and aspects of the physical sciences.

Complimentary online access to the first volume will be available until January 2015.

TABLE OF CONTENTS:

- *What Is Statistics?* Stephen E. Fienberg
- *A Systematic Statistical Approach to Evaluating Evidence from Observational Studies*, David Madigan, Paul E. Stang, Jesse A. Berlin, Martijn Schuemie, J. Marc Overhage, Marc A. Suchard, Bill Dumouchel, Abraham G. Hartzema, Patrick B. Ryan
- *The Role of Statistics in the Discovery of a Higgs Boson*, David A. van Dyk
- *Brain Imaging Analysis*, F. DuBois Bowman
- *Statistics and Climate*, Peter Guttorp
- *Climate Simulators and Climate Projections*, Jonathan Rougier, Michael Goldstein
- *Probabilistic Forecasting*, Tilmann Gneiting, Matthias Katzfuss
- *Bayesian Computational Tools*, Christian P. Robert
- *Bayesian Computation Via Markov Chain Monte Carlo*, Radu V. Craiu, Jeffrey S. Rosenthal
- *Build, Compute, Critique, Repeat: Data Analysis with Latent Variable Models*, David M. Blei
- *Structured Regularizers for High-Dimensional Problems: Statistical and Computational Issues*, Martin J. Wainwright
- *High-Dimensional Statistics with a View Toward Applications in Biology*, Peter Bühlmann, Markus Kalisch, Lukas Meier
- *Next-Generation Statistical Genetics: Modeling, Penalization, and Optimization in High-Dimensional Data*, Kenneth Lange, Jeanette C. Papp, Janet S. Sinsheimer, Eric M. Sobel
- *Breaking Bad: Two Decades of Life-Course Data Analysis in Criminology, Developmental Psychology, and Beyond*, Elena A. Erosheva, Ross L. Matsueda, Donatello Telesca
- *Event History Analysis*, Niels Keiding
- *Statistical Evaluation of Forensic DNA Profile Evidence*, Christopher D. Steele, David J. Balding
- *Using League Table Rankings in Public Policy Formation: Statistical Issues*, Harvey Goldstein
- *Statistical Ecology*, Ruth King
- *Estimating the Number of Species in Microbial Diversity Studies*, John Bunge, Amy Willis, Fiona Walsh
- *Dynamic Treatment Regimes*, Bibhas Chakraborty, Susan A. Murphy
- *Statistics and Related Topics in Single-Molecule Biophysics*, Hong Qian, S.C. Kou
- *Statistics and Quantitative Risk Management for Banking and Insurance*, Paul Embrechts, Marius Hofert

Access this and all other Annual Reviews journals via your institution at www.annualreviews.org.

ANNUAL REVIEWS | Connect With Our Experts

Tel: 800.523.8635 (US/CAN) | Tel: 650.493.4400 | Fax: 650.424.0910 | Email: service@annualreviews.org

