

Estimation of Genetic Correlation via Linkage Disequilibrium Score Regression and Genomic Restricted Maximum Likelihood

Guiyan Ni,^{1,2} Gerhard Moser,^{1,2} Schizophrenia Working Group of the Psychiatric Genomics Consortium, Naomi R. Wray,^{3,4} and S. Hong Lee^{1,2,4,*}

Genetic correlation is a key population parameter that describes the shared genetic architecture of complex traits and diseases. It can be estimated by current state-of-art methods, i.e., linkage disequilibrium score regression (LDSC) and genomic restricted maximum likelihood (GREML). The massively reduced computing burden of LDSC compared to GREML makes it an attractive tool, although the accuracy (i.e., magnitude of standard errors) of LDSC estimates has not been thoroughly studied. In simulation, we show that the accuracy of GREML is generally higher than that of LDSC. When there is genetic heterogeneity between the actual sample and reference data from which LD scores are estimated, the accuracy of LDSC decreases further. In real data analyses estimating the genetic correlation between schizophrenia (SCZ) and body mass index, we show that GREML estimates based on ~150,000 individuals give a higher accuracy than LDSC estimates based on ~400,000 individuals (from combined meta-data). A GREML genomic partitioning analysis reveals that the genetic correlation between SCZ and height is significantly negative for regulatory regions, which whole genome or LDSC approach has less power to detect. We conclude that LDSC estimates should be carefully interpreted as there can be uncertainty about homogeneity among combined meta-datasets. We suggest that any interesting findings from massive LDSC analysis for a large number of complex traits should be followed up, where possible, with more detailed analyses with GREML methods, even if sample sizes are lesser.

Genetic correlation is a key population parameter that describes the shared genetic architecture of complex traits and diseases.^{1–3} The genetic correlation is the additive genetic covariance between two traits scaled by the square root of the product of the genetic variance for each trait (i.e., the geometric mean of the trait variances). The sign of the correlation shows the direction of sharing, and the parameter definition is based on genetic variants across the allelic spectrum. Methods to estimate genetic correlation based on genetic covariance structure are well established for both quantitative and disease traits, e.g., (restricted) maximum likelihood for linear mixed models (LMM).^{4–6} Genetic covariance structure can be derived from phenotypic records using pedigree information in twin or family-based designs.⁷ Recently, genome-wide single-nucleotide polymorphism (SNP) data have been used to construct a genomic relationship matrix for the genetic covariance structure in LMM that captures the contribution of causal variants that are in linkage disequilibrium (LD) with the genotyped SNPs.^{4,8,9} Such estimates assume that the genetic correlation estimated from common SNPs is representative of the parameter that depends on all genetic variants; this seems like a reasonable assumption.

In contrast to the genomic restricted maximum likelihood (GREML) approach, a linkage disequilibrium score regression (LDSC)^{10,11} method does not require individual-level genotype data but instead uses GWAS summary statistics, regressing association test statistics of SNPs on

their LD scores. The LD score of a SNP is the sum of LD r^2 measured with all other SNPs and can be calculated in a reference sample of the same ethnicity when individual genotype data are not available for the GWAS sample, under the assumption that the GWAS sample has been drawn from the same ethnic population as the reference sample used to calculate the LD scores. The method exploits the relationship between association test statistic and LD score expected under polygenicity. Because of this simplicity, and the massively reduced computing burden in terms of memory and time, it is feasible for LDSC to be applied to a large number of multiple traits, e.g., Bulik-Sullivan et al.,¹¹ Zheng et al.,¹² Finucane et al.¹³

Given the attractiveness of LDSC for a massive analysis of many sets of GWAS summary statistics, it has been widely used in the community. However, genetic correlations estimated by LDSC are often reported without caution although the approach is known to be less accurate, compared to GREML.¹¹ In fact, the accuracies of LDSC estimates have not been thoroughly studied.

In this report, we compare both the bias (difference between the simulated true value and estimated value) and accuracy (magnitude of the standard error of an estimate [SE]) between GREML and LDSC for estimation of genetic correlation. We find that both methods show little evidence of bias. However, LDSC is less accurate as reported in Bulik Sullivan et al.,¹¹ with SE at least more than 1.5-fold higher than that of GREML regardless of the number of samples in data used to estimate the genetic correlation.

¹Centre for Population Health Research, School of Health Sciences and Sansom Institute of Health Research, University of South Australia, Adelaide, SA 5000, Australia; ²School of Environmental and Rural Science, University of New England, Armidale, NSW 2351, Australia; ³Institute for Molecular Bioscience, University of Queensland, Brisbane, QLD 4072, Australia; ⁴Queensland Brain Institute, University of Queensland, Brisbane, QLD 4072, Australia

*Correspondence: hong.lee@unisa.edu.au
<https://doi.org/10.1016/j.ajhg.2018.03.021>

© 2018 American Society of Human Genetics.




Figure 1. The Ratio of SE of LDSC Estimate to that of GREML Estimate using Simulated Phenotypes Based on UK Biobank Genotypes

Bars are 95% CI based on 100 replicates. The unit for the number of SNPs is thousands. This result was based on 858K SNPs (after QC) and 10,000 individuals that were randomly selected from UK Biobank. SNPs in each bin were randomly drawn from the 858K SNPs independently. The number of causal SNPs was 10,000 that were randomly selected in each bin. The true simulated value for the genetic correlation was 0.6 and that for the heritability was 0.5 for both traits. Overlap (0%, 10%, and 20%) stands for the percentage of overlapping individuals in the first and second traits.

homogeneity. Furthermore, a GREML genomic partitioning analysis reveals that the genetic correlation between SCZ and height is significantly negative for regulatory regions, which is less obvious by LDSC when using both whole-genome and partitioned estimates of genetic correlation.

In the main methods, we used GREML^{14,15} and LDSC^{10,11} to compare their estimates of genetic correlation using simulated as well as real data. Simulations were based on UK Biobank imputed genotype data (UKBB¹⁶) after stringent quality control (QC) (see [Supplemental Methods](#)). We calculated a ratio of empirical SE and its 95% confidence interval (CI) to assess the accuracy of

When decreasing the number of SNPs, the accuracy of LDSC decreases further. When increasing the degree of genetic heterogeneity between the actual sample and reference data from which LD scores are estimated, the SE of LDSC estimates are up to 3-fold larger than those of the GREML estimates. We also show that GREML is more accurate in genomic partitioning analyses over LDSC or stratified LDSC (sLDSC). In genomic partitioning analyses, the genetic parameters are estimated for genomic subsets defined by user-specified annotations. In analyses of real data, we show that GREML is more accurate and powerful, e.g., GREML estimates based on ~150,000 individuals give a higher accuracy than LDSC estimates based on 400,000 individuals in estimating genetic correlation between schizophrenia (SCZ) and body mass index (BMI) (-0.136 [SE = 0.017] and p value = $4.54E-15$ for GREML versus -0.087 [SE = 0.019] and p value = $4.91E-06$ for LDSC). In these analyses, the GREML estimate is based on UK sample only whereas the LDSC estimate is based on combined meta-datasets among which there is uncertainty about

the methods for each set of simulated data. The 95% CIs of SE were estimated based on the delta method.¹⁷ When estimating genetic correlation using simulated phenotypes based on UKBB genotype data, we found that the estimates were unbiased for both GREML and LDSC ([Figure S1](#)), but the SE of GREML was at least 1.5 times smaller than that of LDSC ([Figure 1](#)). The ratio of the empirical SE from LDSC to GREML was increased up to 3.5-fold when using a smaller number of SNPs ([Figure 1](#)). All values of the ratio were significantly different from 1. It is notable that the SE of GREML estimates showed almost no difference across different numbers of SNPs whereas that of LDSC estimates gradually increased with a smaller number of SNPs ([Figure S2](#)). The ratio was invariant to sample size ([Figure S3](#)). As expected, when using the intercept constrained to zero, LDSC estimates were substantially biased when there were overlapping samples ([Figure S4](#)). We also explored alternative genetic architectures ([Figure S5](#)), which consistently showed that GREML gives a smaller SE than LDSC in any scenario.

Table 1. Correlation between LD Scores Estimated Based on the HapMap3 SNPs using the 1KG CEU Reference Sample and that from Different Target Populations

	Correlation	Nr.SNPs
UKBB ^a	0.946	858,991
UKBB ^b	0.720	123,615 ^c
WTCCC2	0.899	421,035 ^c
GERA	0.661	238,089 ^c

^aUKBB was imputed to the combined data of the 1KG reference and UK10K data.

^bUKBB^r was based on the raw genotype data of UK Biobank data.

^cThe number of SNPs reduced further from the set of the QCed SNPs because of using only SNPs matched with the HapMap3 SNPs used in calculating LD scores.

To explore the stability of the accuracy for both methods, we used two additional genotype datasets without imputation, Wellcome Trust Case Control Consortium 2 (WTCCC2^{18–21}) and genetic epidemiology research on adult health and aging cohort (GERA^{22,23}), which are publicly available (see [Supplemental Methods](#) for detailed data descriptions). We also used UKBB raw (non-imputed) genotype data (UKBB^r). We calculated the correlation between the LD scores for the HapMap3 SNPs estimated based on the 1KG CEU reference sample (see [Web Resources](#)) and those based on in-sample genotype data, i.e., UKBB, WTCCC2, GERA, and UKBB^r dataset ([Table 1](#)). We found that the WTCCC2, GERA, and UKBB^r (raw) genotypes were less similar to the 1KG reference genotypes, compared to the UKBB (imputed) genotypes (noting that UKBB samples had been imputed to the combined data of 1KG reference and UK10K data). [Table 2](#) shows that the SE ratio of LDSC estimate to GREML estimate was higher for WTCCC2, GERA, and UKBB^r than that for UKBB. [Figure 2](#) shows that the accuracy of GREML was consistent across different datasets, whereas that of LDSC was decreased for WTCCC2, GERA, or UKBB^r, compared to UKBB dataset. This was probably due to higher (or lower) correlation between LD scores based on the 1KG reference and the in-sample genotype datasets ([Table 1](#)) which might positively or (negatively) affect the accuracy of LDSC estimates. For WTCCC2, GERA, and UKBB^r data, the SE ratio of LDSC to GREML based on different number of individuals is shown in [Figures S6–S8](#).

Genome partitioning analyses are an emerging tool to estimate the genetic variance and covariance explained by functional categories (e.g., DNase I hypersensitive sites [DHS] and non-DHS²⁴). Currently, genomic partitioning analyses focus on SNP-heritability enrichment analyses, formally testing for enrichment of signal compared to the expectation that the estimates are proportional to the number of SNPs allocated to each annotation. Considering genomic partitioning in cross-disorder analyses is a natural extension to identify regions where genetic correlations between disorders are highest and lowest. Here, we assessed the performance of the methods in the context of genome partitioning analyses using simulated phenotypes

Table 2. The Ratio of SE of LDSC Estimate to That of GREML Estimate using Simulated Phenotypes Based on UKBB, WTCCC2, GERA, and UKBB^r Genotypes in the Scenarios without Overlapping Individuals

	800k	400k	200k	100k
UKBB	1.60 (0.15)	1.70 (0.18)	1.85 (0.25)	2.04 (0.33)
WTCCC2	NA	2.15 (0.31)	2.35 (0.43)	2.68 (0.61)
GERA	NA	NA	2.87 (0.56)	3.31 (1.17)
UKBB ^r	NA	NA	NA	3.74 (0.79)

based on UKBB genotype data. A better LDSC approach to estimate genetic correlation for each category might be sLDSC, stratifying by genomic annotation; however, this method is currently under development (i.e., there is software [see [Web Resources](#)], but there is no published document or paper verifying the method). Nonetheless, since the sLDSC is available to the research community, we applied both LDSC and sLDSC to estimate partitioned genetic correlations for the simulated data ([Supplemental Methods](#)). For genome partitioning analyses, we showed that LDSC estimates of genetic correlation were biased whether using LD scores estimated from the 1KG reference or in-sample data (UKBB) while GREML estimates gave unbiased estimates for each functional category ([Figure 3](#)). sLDSC estimates were unbiased only when using LD scores from the in-sample data, and their SEs are relatively larger than those of GREML or LDSC ([Figure 3](#)). This was probably due to the fact that the different distribution of causal variants and their effects between DHS and non-DHS regions were better captured by an explicit covariance structure fitted in GREML. We also applied the methods to a range of simulation scenarios and found similar results in that GREML performed better than LDSC or sLDSC ([Figure S9](#) and [Table S1](#)), which was consistent with the previous results ([Figures 1](#) and [2](#)). It is notable that in a deliberately severe scenario (e.g., causal variants are simulated only within few kb of a boundary), GREML could give biased estimation of genetic correlation.^{13,24}

While focusing on the accuracy of genetic correlation estimates, there is an important implication for the bias in SNP-heritability estimates for both GREML and LDSC ([Figure S10](#)). When using the WTCCC2, GERA, and UKBB^r data, which were less similar to the 1KG reference genotypes, compared to the UKBB data, LDSC estimates were substantially biased whereas GREML estimates were close to the true value in estimation of SNP heritability ([Figure S10](#)). However, this result is well known and LDSC was not recommended for SNP heritability by the original authors,¹⁰ but rather only for relative enrichment analysis. Despite this, LDSC is widely used for SNP-heritability estimation (because it is quick and simple). Thus, for completeness we include analyses for different scenarios to quantify the properties of the methods. When reducing the number of SNPs, estimated SNP heritabilities from LDSC were consistently unbiased; however, those from GREML were proportionally underestimated ([Figure S11](#)).




Figure 2. Estimated Genetic Correlation with GREML and LDSC (without Constraint to the Intercept) Based on Different Genetic Datasets

Simulation was based on 10,000 individuals that were randomly selected from UKBB, WTCCC2, GERA, and UKBBr (the raw genotype of UKBB), with 858K, 432K, 239K, and 124K SNPs, respectively. Bars are 95% CI based on 100 replicates. Overlap (0%, 10%, and 20%) stands for the percentage of overlapping individuals in the first and second traits. The gray dashed line stands for the true simulated genetic correlation 0.6.

many statistics for LDSC to see how much the SE of estimates could be reduced by increasing the number of samples and number of SNPs. For real data analyses, we obtained theoretical SE to assess the accuracy of the methods. GREML and LDSC estimates for the SNP heritability were 0.192 (SE 0.004) and 0.280 (SE 0.016) for SCZ and 0.184 (SE 0.004) and 0.255 (SE 0.014) for BMI. The notable difference between GREML and LDSC was probably because of a relatively small number of SNPs (500K) that might result in underestimated GREML SNP heritability (see [Figure S11](#)). This is one of the caveats of using GREML with real data that usually comprise multiple cohorts genotyped on different platforms, such that, even with imputation, the overlapping set of SNPs imputed with high confidence may be limited. The estimated genetic correlation for GREML and LDSC was -0.136

(SE 0.017) and -0.173 (SE 0.031). This indicated that the GREML estimate was 3.5 and 1.8 times more precise than LDSC estimates for the SNP heritability and genetic correlation, respectively. For LDSC, we also considered using additional GWAS summary statistics from publicly available resources.^{25,26} The sample sizes used for additional LDSC analyses (LDSC-meta) are summarized in [Table 3](#). The estimated SNP heritability was 0.259 (SE 0.019) for SCZ and 0.121 (SE 0.007) for BMI, and the estimated genetic correlation was -0.087 (SE 0.019). Although sample size was increased 2.7-fold, the SE of LDSC estimate was not smaller than that for GREML estimate (SE = 0.017 versus 0.019, and p value = $4.54E-15$ versus $4.91E-06$ for GREML versus LDSC) ([Table 3](#)). It should be noted that GREML estimates used a homogeneous population (within UK and after stringent QC excluding population outliers) whereas LDSC-meta1 and -meta2 were based on

When using non-HapMap3 SNPs, LDSC estimates were consistently biased ([Figure S12](#)) and less accurate, compared to GREML estimates ([Figures S13](#) and [S14](#)), which probably explains why LDSC is implemented using only HapMap3 SNPs. Although the genetic correlation is robust to such biasedness,^{4,11} SNP heritability itself should be carefully interpreted for both GREML and LDSC. We also noted that LDSC and sLDSC estimates for SNP heritability were biased in the genome partitioning analysis ([Figure S15](#)) although the estimated enrichment was close to the true value when using sLDSC and in-sample LD scores ([Figure S15](#)).

We used real phenotype and individual genotype data from the Psychiatric Genomics Consortium (PGC) and UKBB to estimate genetic variance and covariance between SCZ and BMI using LDSC and GREML ([Table 3](#) and [Figure S16](#)). We also used publicly available GWAS sum-




Figure 3. Estimated Genetic Correlation of Simulated Data Based on a Genomic Partitioning Model

Simulation was based on 10,000 individuals that were randomly selected from UKBB with 858K SNP. Based on Gusev et al.,²⁴ the 858K SNPs across the genome were stratified as two categories: DHS (194K SNPs with 2,268 causal SNPs) and non-DHS (664K SNPs with 7,732 causal SNPs). The genetic correlation for the simulated phenotypes between the first and second traits was 0.6 and -0.6 in DHS and non-DHS region, respectively. Bars are 95% CI based on 100 replicates. LDSC-CEU: Using LD-scores estimated from 1KG reference data. LDSC-OWN: Using LD-scores estimated from UKBB. sLDSC-CEU: Using stratified LD-scores estimated from 1KG reference data. sLDSC-OWN: Using stratified LD-scores estimated from UKBB. The presented results were based on 0% overlapping samples between the first and second traits and those based on other scenarios (e.g., 10% and 20%) are presented in [Table S1](#).

and [S20](#)), and show that the estimates were similar between LDSC and GREML. However, GREML had a lower p value (0.0028 in [Figure 4](#)) than LDSC using LD scores from the 1KG reference data (p value = 0.04) or using LD scores from the in-sample data (p value = 0.007). We note that current sLDSC software does not provide a SE of estimated partitioned genetic correlation for each category;

combined meta-datasets consisting of ~80 different studies for which there is much more uncertainty about homogeneity than when using a single study cohort such as UKBB. The large difference of the estimates between LDSC and LDSC-meta1 (or -meta2) was probably due to the fact that heterogeneity among the 80 different studies resulted in underestimation of the common genetic variance and covariance, and that the difference of LD scores between the target and 1KG reference data would bias the LDSC estimates as shown in [Figure S10](#). We also analyzed height data²⁷ and found a similar pattern in that GREML estimates were more accurate than LDSC estimates whether using the same data or using additional GWAS summary statistics for LDSC ([Figure S17](#) and [Table S2](#)).

In the real data analyses, we carried out a functional category analysis partitioning the genome into regulatory, DHS, intronic, and intergenic regions using GREML ([Figure 4](#) for SCZ/height and [Figure S18](#) for SCZ/BMI). For SCZ and height, the genetic correlation for the regulatory region was negative and significantly different from 0 (p value = 0.0028; [Figure 4](#)). We also compared the results with the LDSC genetic correlation estimation ([Figures S19](#)

therefore, we did not attempt using the software for the real data analysis. For SNP-heritability estimation, the SE of the estimate for each category was much lower for GREML than sLDSC, ranging from 2.2- to 5.9-fold ([Table S3](#)).

LDSC and GREML are the methods that have been widely used in estimating genetic correlation, shedding light on the shared genetic architecture of complex traits, based on genome-wide SNPs. Two critical parameters for assessing methods are bias (whether the estimates over replicated analyses differ from the true value) and accuracy (reflected by the standard error of the estimate). Although the property of the accuracy of GREML has been thoroughly studied and tested,^{29,30} that of LDSC has not been sufficiently investigated. In this report, we compare the accuracy of GREML and LDSC estimates based on various scenarios using simulated as well as real datasets, and draw simple but useful guidelines ([Box 1](#)).

Both GREML and LDSC are methods that aim to estimate the same genetic correlation parameter based on genetic variants across the allelic spectrum as defined earlier and the definition is invariant across the methods. The estimates from both GREML and LDSC are valid if all

Table 3. Heritability and Genetic Correlation Based on Different Datasets

Method	#SNPs	Data	# Individuals		h ² BMI		h ² SCZ (Liability Scale)		Genetic Correlation		
			Mean	SD	Estimate	SE	Estimate	SE	Estimate	SE	p
GREML	518,992	UKBB+SCZ(qced)	152,961	–	0.184	3.80E–03	0.192	4.39E–03	–0.136	1.74E–02	4.54E–15
LDSC	516,519	UKBB+SCZ(qced)	151,262	1,432.7	0.255	1.38E–02	0.280	1.63E–02	–0.173	3.08E–02	1.91E–08
LDSC-meta1	477,163	UKBB+GIANT+PGCSCZ	422,499	20,226.0	0.111	8.10E–03	0.259	1.28E–02	–0.091	2.44E–02	1.95E–04
LDSC-meta2	1,011,748	UKBB+GIANT+PGCSCZ	414,707	32,697.8	0.121	6.50E–03	0.261	1.03E–02	–0.087	1.90E–02	4.91E–06

GREML: Analysis was based on quality controlled genetic data for BMI (from UK Biobank with 111,019 individuals and 518,992 SNPs) and schizophrenia (from PGC with 41,630 individuals and 518,992 SNPs).

LDSC: The datasets used in LDSC were the same as in GREML.

LDSC-meta1: GWAS summary statistics for BMI were based on meta-analyzed GWAS results of UKBB individual-level genetic data (with 111,019 individuals and 518,992 SNPs) and of GIANT (245,051 individuals and 477,163 SNPs). For SCZ, the GWAS summary statistics from the full PGC sample based on 77,096 individuals were used.

LDSC-meta2: The datasets used in LDSC-meta2 were the same as in LDSC-meta1 except the increased number of SNPs (1,011,748) with which its performance was to check.

Mean and SD of #individuals: Due to different call rates of each SNP, number of individuals for each SNP used in GWAS were different.

required assumptions are met. GREML estimates variance/covariance components based on genetic covariance structure estimated from available (in-sample) individual genotypes, whereas LDSC estimates variance/covariance components based on association test statistics corrected for LD structure inferred from the markers in the reference panel (e.g., 1KG of the same ethnicity). The underlying assumption is that the samples generating the GWAS summary statistics are drawn from the same population as the samples generating the LDSC statistics, but here we showed that there can be LD-structure (LD-scores) differences between in-sample and reference data, which impacts parameter estimations (Tables 1 and 2 and Figure S10).

The reduced computing burden of LDSC over GREML makes it the method of choice for generating a quick overview of the genetic relationship between disorders (Table S4). However, our results suggest that important associations could be overlooked. For example, Bulik-Sullivan et al.¹¹ reported a negative genetic correlation between BMI and SCZ estimated by LDSC (estimate = –0.095, SE = 0.025 with p value = 1.75E–4), which was not significant after Bonferroni correction for the multiple testing. Because of the limited power from LDSC analysis, the shared genetic architecture between BMI and SCZ, perhaps, has had less attention than it is due. We confirmed the negative genetic correlation between BMI and SCZ with a greater confidence (estimate = –0.136, p value = 4.54E–15) using GREML. A second example is in analyses investigating the shared genetic architecture between height and SCZ, in which epidemiological evidence points to a negative association,³¹ supported by genetic analyses.³² However, there was no evidence of genetic correlation between height and SCZ in whole-genome level analyses of Bulik-Sullivan et al.¹¹ (estimate = –0.002, SE = 0.022). We used a GREML genomic partitioning analysis and found a significant negative genetic correlation between height and SCZ for the regulatory region (Figure 4). It was noted that the regulatory region was highly enriched for height (estimate = 0.094, p value = 7.60E–92 in Table S3), which

intuitively supports a significant genetic correlation with SCZ for the region. As shown in Figures 3 and S15, the GREML estimate was closer to the true values with a lower SE than LDSC or sLDSC estimate in simulated data. For the real data analyses (Table S3), GREML had more accurate SNP-heritability estimates (lower SE) than sLDSC. Moreover, the sum of each category matched well with the estimate of the whole genome for GREML whereas this was not the case for sLDSC (Tables S3).

Here we focused on genetic correlation estimates and did not consider a number of alternative approaches that have been explored in detail for estimation of SNP heritability, e.g., LDK approach,³³ weighted genomic relationship matrix,³⁴ MAF stratified,²⁹ and LD-MAF stratified approaches.³⁵ It was beyond the scope of our study to assess whether biasedness and accuracy can be improved with these methods, although a general observation is that biases in SNP-heritability estimation can “cancel” in estimates of genetic correlations, as biases impact both the numerator and denominator of the genetic correlation quotient.^{4,11} We note that while under review, two new methods to estimate stratified genetic correlations via GWAS summary statistics^{36,37} have been published as alternatives to sLDSC. Those approaches also need external reference samples to infer LD structure in the actual sample, implying the same problem as for LDSC (4 and 5 in Box 1). However, to partially address this problem, one method³⁶ achieves smaller standard errors than sLDSC through a block diagonalization of the LD matrix. A further study is needed to make explicit comparisons with GREML.

In conclusion, LDSC may be the best tool for a massive analysis of multiple sets of GWAS summary statistics in estimating genetic correlation between complex traits, because of its low computing burden and because summary statistics may be available for much larger sample sizes than those with individual genotype data. However, LDSC estimates should be carefully interpreted, considering the summary points (Box 1). Any interesting findings from LDSC analyses should be followed up, where




Figure 4. Genetic Correlation between SCZ and Height and Heritability Based on SNPs in Partitioned Genomic Regions Estimated with GREML

A joint model was applied by fitting four genomic relationship matrices simultaneously, each estimated based on the set of SNPs belong to each of the functional categories (regulatory, intron, intergene, and DHS). The bars are standard errors. *p* value for the estimate significantly different from 0 was 0.0028, 0.52, 0.91, and 0.67 for regulatory, intronic, intergenic, and DHS region, respectively.

possible, with more detailed analyses using individual genotype data and with GREML methods, even though sample sizes with individual genotype data may be smaller.

Supplemental Data

Supplemental Data include 23 figures, 4 tables, Supplemental Methods, and Consortia Members and Affiliations and can be

found with this article online at <https://doi.org/10.1016/j.ajhg.2018.03.021>.

Consortia

The members of the Schizophrenia Working Group of the Psychiatric Genomics Consortium are Stephan Ripke, Benjamin M. Neale, Aiden Corvin, James T.R. Walters, Kai-How Farh, Peter A. Holmans, Phil Lee, Brendan Bulik-Sullivan, David A. Collier, Hailiang Huang, Tune H. Pers, Ingrid Agartz, Esben Agerbo, Margot Albus, Madeline Alexander, Farooq Amin, Silviu A. Bacanu, Martin Begemann, Richard A. Belliveau Jr, Judit Bene, Sarah E. Bergen, Elizabeth Bevilacqua, Tim B. Bigdeli, Donald W. Black, Richard Bruggeman, Nancy G. Buccola, Randy L. Buckner, William Byerley, Wiepke Cahn, Guiqing Cai, Dominique Champion, Rita M. Cantor, Vaughan J. Carr, Noa Carrera, Stanley V. Catts, Kimberly D. Chambert, Raymond C.K. Chan, Ronald Y.L. Chen, Eric Y. H. Chen, Wei Cheng, Eric F.C. Cheung, Siow Ann Chong, C. Robert Cloninger, David Cohen, Nadine Cohen, Paul Cormican, Nick Craddock, James J. Crowley, David Curtis, Michael Davidson, Kenneth L. Davis, Franziska Degenhardt, Jurgen Del Favero, Ditte Demontis, Dimitris Dikeos, Timothy Dinan, Srdjan Djurovic, Gary Donohoe, Elodie Drapeau, Jubao Duan, Frank Dudbridge, Naser Durmishi, Peter Eichhammer, Johan Eriksson, Valentina Escott-Price, Laurent Essioux, Ayman H. Fanous, Martilias S. Farrell, Josef Frank, Lude Franke, Robert Freedman, Nelson B. Freimer, Marion Friedl, Joseph I. Friedman, Menachem Fromer, Giulio Genovese, Lyudmila Georgieva, Ina Giegling, Paola Giusti-Rodríguez, Stephanie Godard, Jacqueline I. Goldstein, Vera Golimbet, Srihari Gopal, Jacob Gratten, Lieuwe de Haan, Christian Hammer, Marian L. Hamshere, Mark Hansen, Thomas Hansen, Vahram Haroutunian, Annette M. Hartmann, Frans A. Henskens, Stefan Herms, Joel N. Hirschhorn, Per Hoffmann, Andrea Hofman, Mads V. Hollegaard, David M. Hougaard, Masashi Ikeda, Inge Joa, Antonio Juliá, René S. Kahn, Luba Kalaydjieva, Sena Karachanak-Yankova, Juha Karjalainen, David Kavanagh, Matthew C. Keller, James L. Kennedy, Andrey Khrunin, Yunjung Kim, Janis Klovins, James A. Knowles, Bettina Konte, Vaidutis Kucinskas, Zita Ausrele Kucinskiene, Hana Kuzelova-Ptackova, Anna K. Kähler, Claudine Laurent, Jimmy Lee Chee Keong, Sophie E. Legge, Bernard Lerer, Miaoxin Li, Tao Li, Kung-Yee Liang, Jeffrey Lieberman, Svetlana Limborska, Carmel M. Loughland, Jan Lubinski, Jouko Lönnqvist, Milan Macek Jr, Patrik K.E. Magnusson, Brion S. Maher, Wolfgang Maier, Jacques Mallet, Sara Marsal, Manuel Mattheisen, Morten Mattingdsda, Robert W. McCarley, Colm McDonald, Andrew M. McIntosh, Sandra Meier, Carin J. Meijer, Bela Melegh, Ingrid Melle, Raquelle I. Mesholam-Gately, Andres Metspalu, Patricia T. Michie, Lili Milani, Vihra Milanova, Younes Mokrab, Derek W. Morris, Ole Mors, Kieran C. Murphy, Robin M. Murray, Inez Myin-Germeys, Bertram Müller-Myhsok, Mari Nelis, Igor Nenadic, Deborah A. Nertney, Gerald Nestadt, Kristin K. Nicodemus, Liene Nikitina-Zake, Laura Nisenbaum, Annelie Nordin, Eadbhard O'Callaghan, Colm O'Dushlaine, F. Anthony O'Neill, Sang-Yun Oh, Ann Olin, Line Olsen, Jim Van Os, Psychosis Endophenotypes International Consortium, Christos Pantelis, George N. Papadimitriou, Sergi Papio, Elena Parkhomenko, Michele T. Pato, Tiina Paunio, Milica Pejovic-Milovancevic, Diana O. Perkins, Olli Pietiläinen, Jonathan Pimm, Andrew J. Pocklington, John Powell, Alkes Price, Ann E. Pulver, Shaun M. Purcell, Digby Quedsted, Henrik B. Rasmussen, Abraham Reichenberg, Mark A. Reimers, Alexander L. Richards, Joshua L. Roffman, Panos Roussos, Douglas M. Ruderfer, Veikko Salomaa, Alan R. Sanders,

Box 1. Summary Points

1. GREML and LDSC can both provide unbiased estimates of the genetic correlation between two traits. GREML requires individual-level genotype data, while LDSC requires only association summary statistics and LD scores per SNP. If LD scores have been calculated from the same sample as the association statistics, then GREML and LDSC provide similar estimates of the genetic correlation. However, in practice LD scores are estimated from external reference samples of the same broad ethnicity, which can lead to bias in the estimates (Figures S21 and S22). As a rule of thumb, when LDSC and GREML estimates are dissimilar, we recommend reporting the estimate with a lower SE. The theoretical SE of the estimates is a reliable indicator to determine the better estimator, which agrees well with the empirical SE (from simulation replicates) (Figure S23).
2. When combining multiple data sets to estimate genetic correlations between multiple traits, it is possible, in practice, that the number of SNPs remaining after QC is relatively small. When the number of available SNPs is small, the SE of LDSC estimates for genetic correlation can be increased relatively more, compared to that of GREML estimates (Figure S2).
3. SNP heritability has a different property, compared to genetic correlation since the latter is robust to biased estimation of genetic variance and covariance (presumably the biases occur in the numerator and denominator and hence approximately cancel out).^{4,11} Especially when using a small number of SNPs (<500K) for GREML or when using multiple meta-data sets for LDSC, estimated SNP heritability itself should be reported with caution as both methods can give biased estimates.
4. When using a study cohort, it is desirable to measure heterogeneity between the cohort and 1KG reference data (e.g., measuring the correlation between LD scores estimated based on the cohort and 1KG reference data as in Table 1). If the correlation is not close to 1, LDSC estimates should be carefully interpreted. We recommend that when GWAS summary statistics are provided, cohort-specific LD scores are provided also. It is also warranted that an optimal approach to meta-analyze LD scores across multiple cohorts should be developed to improve LDSC performance.²⁸
5. When using extensive metadata that possibly include heterogeneous sources, there are two problems. Firstly, the LD scores estimated from reference samples such as 1KG reference may be a poor representation of the LD scores of the heterogeneous metadata, such that the accuracy of LDSC decreases. Second, the distribution of causal variants and pleiotropic effects may be different between heterogeneous sources such that the estimates can be biased (capturing only common effects between heterogeneous sources). This implies that LDSC estimates should be reported with caution when using extensive metadata sets (Table 3).
6. One of advantages of having access to individual-level genotype data comes when more detailed analyses are required, such as genomic partitioning analyses. As shown in Figure 4, a GREML genomic partitioning analysis reveals a significant negative genetic correlation between SCZ and height for the regulatory region, which genome-wide GREML or LDSC approach has less power to detect.

Ulrich Schall, Christian R. Schubert, Thomas G. Schulze, Sibylle G. Schwab, Edward M. Scolnick, Rodney J. Scott, Larry J. Seidman, Jianxin Shi, Engilbert Sigurdsson, Teimuraz Silagadze, Jeremy M. Silverman, Kang Sim, Petr Slominsky, Jordan W. Smoller, Hon-Cheong So, Chris C.A. Spencer, Eli A. Stah, Hreinn Stefansson, Stacy Steinberg, Elisabeth Stogmann, Richard E. Straub, Eric Strengman, Jana Strohmaier, T. Scott Stroup, Mythily Subramaniam, Jaana Suvisaari, Dragan M. Svrakic, Jin P. Szatkiewicz, Erik Söderman, Srinivas Thirumalai, Draga Toncheva, Sarah Tosato, Juha Veijola, John Waddington, Dermot Walsh, Dai Wang, Qiang Wang, Bradley T. Webb, Mark Weiser, Dieter B. Wildenauer, Nigel M. Williams, Stephanie Williams, Stephanie H. Witt, Aaron R. Wolen, Emily H.M. Wong, Brandon K. Wormley, Hualin Simon Xi, Clement C. Zai, Xuebin Zheng, Fritz Zimprich, Kari Stefansson, Peter M. Visscher, Wellcome Trust Case-Control Consortium, Rolf Adolfsson, Ole A. Andreassen, Douglas H.R. Blackwood, Elvira Bramon, Joseph D. Buxbaum, Anders D. Børghlum, Sven Cichon, Ariel Darvasi, Enrico Domenici, Hannelore Ehrenreich, Tõnu Esko, Pablo V. Gejman, Michael Gill, Hugh Gurling, Christina M. Hultman, Nakao Iwata, Assen V. Jablensky, Erik G. Jönsson, Kenneth

S. Kendler, George Kirov, Jo Knight, Todd Lencz, Douglas F. Levinson, Qingqin S. Li, Jianjun Liu, Anil K. Malhotra, Steven A. McCauley, Andrew McQuillin, Jennifer L. Moran, Preben B. Mortensen, Bryan J. Mowry, Markus M. Nöthen, Roel A. Ophoff, Michael J. Owen, Aarno Palotie, Carlos N. Pato, Tracey L. Petryshen, Danielle Posthuma, Marcella Rietsche, Brien P. Riley, Dan Rujescu, Pak C. Sham, Pamela Sklar, David St Clair, Daniel R. Weinberger, Jens R. Wendland, Thomas Werge, Mark J. Daly, Patrick F. Sullivan, and Michael C. O'Donovan.

Acknowledgments

This research is supported by the Australian National Health and Medical Research Council (1080157, 1087889) and the Australian Research Council (DP160102126, FT160100229). This research has been conducted using the UK Biobank Resource. UK Biobank Research Ethics Committee (REC) approval number is 11/NW/0382. Our reference number approved by UK Biobank is 14575. GERA data came from a grant, the Resource for Genetic

Epidemiology Research in Adult Health and Aging (RC2 AG033067; Schaefer and Risch, PIs) awarded to the Kaiser Permanente Research Program on Genes, Environment, and Health (RPGEH) and the UCSF Institute for Human Genetics. The RPGEH was supported by grants from the Robert Wood Johnson Foundation, the Wayne and Gladys Valley Foundation, the Ellison Medical Foundation, Kaiser Permanente Northern California, and the Kaiser Permanente National and Northern California Community Benefit Programs. The RPGEH and the Resource for Genetic Epidemiology Research in Adult Health and Aging are described in the GERA website (see [Web Resources](#)). This study makes use of data generated by the Wellcome Trust Case-Control Consortium. A full list of the investigators who contributed to the generation of the WTCCC data is available online. Funding for the WTCCC project was provided by the Wellcome Trust under awards 076113, 085475, and 090355.

Received: October 4, 2017

Accepted: March 20, 2018

Published: May 10, 2018

Web Resources

GERA, https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=phs000674.v2.p2

GIANT GWAS results, https://portals.broadinstitute.org/collaboration/giant/index.php/GIANT_consortium_data_files

LD scores, <https://data.broadinstitute.org/alkesgroup/LDSCORE/LDSC>, <https://github.com/bulik/ldsc>

MTG2, <https://sites.google.com/site/honglee0707/mtg2>

Psychiatric Genomics Consortium, <http://www.med.unc.edu/pgc/>

UK Biobank, <http://www.ukbiobank.ac.uk>

WTCCC2, <http://www.wtccc.org.uk/cc2>

References

- Mehta, D., Tropf, F.C., Gratten, J., Bakshi, A., Zhu, Z., Bacanu, S.-A., Hemani, G., Magnusson, P.K.E., Barban, N., Esko, T., et al.; Schizophrenia Working Group of the Psychiatric Genomics Consortium, LifeLines Cohort Study, and TwinsUK (2016). Evidence for genetic overlap between schizophrenia and age at first birth in women. *JAMA Psychiatry* 73, 497–505.
- Lee, S.H., Byrne, E.M., Hultman, C.M., Kähler, A., Vinkhuyzen, A.A.E., Ripke, S., Andreassen, O.A., Frisell, T., Gusev, A., Hu, X., et al.; Schizophrenia Working Group of the Psychiatric Genomics Consortium and Rheumatoid Arthritis Consortium International; Schizophrenia Working Group of the Psychiatric Genomics Consortium Authors; Schizophrenia Working Group of the Psychiatric Genomics Consortium Collaborators; Rheumatoid Arthritis Consortium International Authors; and Rheumatoid Arthritis Consortium International Collaborators (2015). New data and an old puzzle: the negative association between schizophrenia and rheumatoid arthritis. *Int. J. Epidemiol.* 44, 1706–1721.
- Lee, S.H., DeCandia, T.R., Ripke, S., Yang, J., Sullivan, P.F., Goddard, M.E., Keller, M.C., Visscher, P.M., Wray, N.R.; Schizophrenia Psychiatric Genome-Wide Association Study Consortium (PGC-SCZ); International Schizophrenia Consortium (ISC); and Molecular Genetics of Schizophrenia Collaboration (MGS) (2012). Estimating the proportion of variation in susceptibility to schizophrenia captured by common SNPs. *Nat. Genet.* 44, 247–250.
- Lee, S.H., Yang, J., Goddard, M.E., Visscher, P.M., and Wray, N.R. (2012). Estimation of pleiotropy between complex diseases using single-nucleotide polymorphism-derived genomic relationships and restricted maximum likelihood. *Bioinformatics* 28, 2540–2542.
- Harville, D.A. (1977). Maximum likelihood approaches to variance component estimation and to related problems. *J. Am. Stat. Assoc.* 72, 320–338.
- Patterson, H.D., and Thompson, R. (1971). Recovery of interblock information when block sizes are unequal. *Biometrika* 58, 545–554.
- Neale, M., and Cardon, L. (2013). *Methodology for Genetic Studies of Twins and Families* (Springer Science & Business Media).
- VanRaden, P.M. (2008). Efficient methods to compute genomic predictions. *J. Dairy Sci.* 91, 4414–4423.
- Yang, J., Lee, S.H., Goddard, M.E., and Visscher, P.M. (2011). GCTA: a tool for genome-wide complex trait analysis. *Am. J. Hum. Genet.* 88, 76–82.
- Bulik-Sullivan, B.K., Loh, P.R., Finucane, H.K., Ripke, S., Yang, J., Patterson, N., Daly, M.J., Price, A.L., Neale, B.M.; and Schizophrenia Working Group of the Psychiatric Genomics Consortium (2015). LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat. Genet.* 47, 291–295.
- Bulik-Sullivan, B., Finucane, H.K., Anttila, V., Gusev, A., Day, F.R., Loh, P.R., Duncan, L., Perry, J.R., Patterson, N., Robinson, E.B., et al.; ReproGen Consortium; Psychiatric Genomics Consortium; and Genetic Consortium for Anorexia Nervosa of the Wellcome Trust Case Control Consortium 3 (2015). An atlas of genetic correlations across human diseases and traits. *Nat. Genet.* 47, 1236–1241.
- Zheng, J., Erzurumluoglu, A.M., Elsworth, B.L., Kemp, J.P., Howe, L., Haycock, P.C., Hemani, G., Tansey, K., Laurin, C., Pourcain, B.S., et al.; Early Genetics and Lifecourse Epidemiology (EAGLE) Eczema Consortium (2017). LD Hub: a centralized database and web interface to perform LD score regression that maximizes the potential of summary level GWAS data for SNP heritability and genetic correlation analysis. *Bioinformatics* 33, 272–279.
- Finucane, H.K., Bulik-Sullivan, B., Gusev, A., Trynka, G., Reshef, Y., Loh, P.R., Anttila, V., Xu, H., Zang, C., Farh, K., et al.; ReproGen Consortium; Schizophrenia Working Group of the Psychiatric Genomics Consortium; and RACI Consortium (2015). Partitioning heritability by functional annotation using genome-wide association summary statistics. *Nat. Genet.* 47, 1228–1235.
- Lee, S.H., and van der Werf, J.H. (2016). MTG2: an efficient algorithm for multivariate linear mixed model analysis based on genomic information. *Bioinformatics* 32, 1420–1422.
- Maier, R., Moser, G., Chen, G.-B., Ripke, S., Coryell, W., Potash, J.B., Scheftner, W.A., Shi, J., Weissman, M.M., Hultman, C.M., et al.; Cross-Disorder Working Group of the Psychiatric Genomics Consortium (2015). Joint analysis of psychiatric disorders increases accuracy of risk prediction for schizophrenia, bipolar disorder, and major depressive disorder. *Am. J. Hum. Genet.* 96, 283–294.
- Collins, R. (2012). What makes UK Biobank special? *Lancet* 379, 1173–1174.

17. Lynch, M., and Walsh, B. (1998). *Genetics and Analysis of Quantitative Traits* (MA: Sinauer Sunderland).
18. Sawcer, S., Hellenthal, G., Pirinen, M., Spencer, C.C., Patshopoulos, N.A., Moutsianas, L., Dilthey, A., Su, Z., Freeman, C., Hunt, S.E., et al.; International Multiple Sclerosis Genetics Consortium; and Wellcome Trust Case Control Consortium 2 (2011). Genetic risk and a primary role for cell-mediated immune mechanisms in multiple sclerosis. *Nature* *476*, 214–219.
19. Mells, G.F., Floyd, J.A.B., Morley, K.I., Cordell, H.J., Franklin, C.S., Shin, S.-Y., Heneghan, M.A., Neuberger, J.M., Donaldson, P.T., Day, D.B., et al.; UK PBC Consortium; and Wellcome Trust Case Control Consortium 3 (2011). Genome-wide association study identifies 12 new susceptibility loci for primary biliary cirrhosis. *Nat. Genet.* *43*, 329–332.
20. Bellenguez, C., Bevan, S., Gschwendtner, A., Spencer, C.C., Burgess, A.I., Pirinen, M., Jackson, C.A., Traylor, M., Strange, A., Su, Z., et al.; International Stroke Genetics Consortium (ISGC); and Wellcome Trust Case Control Consortium 2 (WTCCC2) (2012). Genome-wide association study identifies a variant in HDAC9 associated with large vessel ischemic stroke. *Nat. Genet.* *44*, 328–333.
21. Tsoi, L.C., Spain, S.L., Knight, J., Ellinghaus, E., Stuart, P.E., Capon, F., Ding, J., Li, Y., Tejasvi, T., Gudjonsson, J.E., et al.; Collaborative Association Study of Psoriasis (CASP); Genetic Analysis of Psoriasis Consortium; Psoriasis Association Genetics Extension; and Wellcome Trust Case Control Consortium 2 (2012). Identification of 15 new psoriasis susceptibility loci highlights the role of innate immunity. *Nat. Genet.* *44*, 1341–1348.
22. Banda, Y., Kvale, M.N., Hoffmann, T.J., Hesselson, S.E., Ranatunga, D., Tang, H., Sabatti, C., Croen, L.A., Dispensa, B.P., Henderson, M., et al. (2015). Characterizing race/ethnicity and genetic ancestry for 100,000 subjects in the genetic epidemiology research on adult health and aging (GERA) cohort. *Genetics* *200*, 1285–1295.
23. Lee, S.H., Weerasinghe, W.M.S.P., Wray, N.R., Goddard, M.E., and van der Werf, J.H.J. (2017). Using information of relatives in genomic prediction to apply effective stratified medicine. *Sci. Rep.* *7*, 42091.
24. Gusev, A., Lee, S.H., Trynka, G., Finucane, H., Vilhjálmsson, B.J., Xu, H., Zang, C., Ripke, S., Bulik-Sullivan, B., Stahl, E., et al.; Schizophrenia Working Group of the Psychiatric Genomics Consortium; SWE-SCZ Consortium; Schizophrenia Working Group of the Psychiatric Genomics Consortium; and SWE-SCZ Consortium (2014). Partitioning heritability of regulatory and cell-type-specific variants across 11 common diseases. *Am. J. Hum. Genet.* *95*, 535–552.
25. Schizophrenia Working Group of the Psychiatric Genomics Consortium (2014). Biological insights from 108 schizophrenia-associated genetic loci. *Nature* *511*, 421–427.
26. Locke, A.E., Kahali, B., Berndt, S.I., Justice, A.E., Pers, T.H., Day, F.R., Powell, C., Vedantam, S., Buchkovich, M.L., Yang, J., et al.; LifeLines Cohort Study; ADIPOGen Consortium; AGEN-BMI Working Group; CARDIOGRAMplusC4D Consortium; CKDGen Consortium; GLGC; ICBP; MAGIC Investigators; MuTHER Consortium; MIGen Consortium; PAGE Consortium; ReproGen Consortium; GENIE Consortium; and International Endogene Consortium (2015). Genetic studies of body mass index yield new insights for obesity biology. *Nature* *518*, 197–206.
27. Wood, A.R., Esko, T., Yang, J., Vedantam, S., Pers, T.H., Gustafsson, S., Chu, A.Y., Estrada, K., Luan, J., Kutalik, Z., et al.; Electronic Medical Records and Genomics (eMEMERGE) Consortium; MIGen Consortium; PAGE Consortium; and LifeLines Cohort Study (2014). Defining the role of common variation in the genomic and biological architecture of adult human height. *Nat. Genet.* *46*, 1173–1186.
28. Pasaniuc, B., and Price, A.L. (2017). Dissecting the genetics of complex traits using summary association statistics. *Nat. Rev. Genet.* *18*, 117–127.
29. Lee, S.H., Yang, J., Chen, G.B., Ripke, S., Stahl, E.A., Hultman, C.M., Sklar, P., Visscher, P.M., Sullivan, P.F., Goddard, M.E., and Wray, N.R. (2013). Estimation of SNP heritability from dense genotype data. *Am. J. Hum. Genet.* *93*, 1151–1155.
30. Speed, D., Hemani, G., Johnson, M.R., and Balding, D.J. (2012). Improved heritability estimation from genome-wide SNPs. *Am. J. Hum. Genet.* *91*, 1011–1021.
31. Gunnell, D., Harrison, G., Whitley, E., Lewis, G., Tynelius, P., and Rasmussen, F. (2005). The association of fetal and childhood growth with risk of schizophrenia. Cohort study of 720,000 Swedish men and women. *Schizophr. Res.* *79*, 315–322.
32. Bacanu, S.A., Chen, X., and Kendler, K.S. (2013). The genetic overlap between schizophrenia and height. *Schizophr. Res.* *151*, 226–228.
33. Speed, D., Cai, N., Johnson, M.R., Nejentsev, S., Balding, D.J.; and UCLEB Consortium (2017). Reevaluation of SNP heritability in complex human traits. *Nat. Genet.* *49*, 986–992.
34. de Los Campos, G., Vazquez, A.I., Fernando, R., Klimentidis, Y.C., and Sorensen, D. (2013). Prediction of complex human traits using the genomic best linear unbiased predictor. *PLoS Genet.* *9*, e1003608–e1003608.
35. Yang, J., Bakshi, A., Zhu, Z., Hemani, G., Vinkhuyzen, A.A.E., Lee, S.H., Robinson, M.R., Perry, J.R.B., Nolte, I.M., van Vliet-Ostapchouk, J.V., et al.; LifeLines Cohort Study (2015). Genetic variance estimation with imputed variants finds negligible missing heritability for human height and body mass index. *Nat. Genet.* *47*, 1114–1120.
36. Shi, H., Mancuso, N., Spendlove, S., and Pasaniuc, B. (2017). Local genetic correlation gives insights into the shared genetic architecture of complex traits. *Am. J. Hum. Genet.* *101*, 737–751.
37. Lu, Q., Li, B., Ou, D., Erlendsdottir, M., Powles, R.L., Jiang, T., Hu, Y., Chang, D., Jin, C., Dai, W., et al. (2017). A powerful approach to estimating annotation-stratified genetic covariance via GWAS summary statistics. *Am. J. Hum. Genet.* *101*, 939–964.