

Multi-trait Genomic Selection Methods for Crop Improvement

Saba Moeinizade*, Aaron Kusmec†, Guiping Hu*,¹, Lizhi Wang* and Patrick S. Schnable†

*Department of Industrial and Manufacturing Systems Engineering, Iowa State University, †Department of Agronomy, Iowa State University

ABSTRACT Plant breeders make selection decisions based on multiple traits, such as yield, plant height, flowering time, and disease resistance. A commonly used approach in multi-trait genomic selection is index selection, which assigns weights to different traits relative to their economic importance. However, classical index selection only optimizes genetic gain in the next generation, requires some experimentation to find weights that lead to desired outcomes, and has difficulty optimizing non-linear breeding objectives. Multi-objective optimization has also been used to identify the Pareto frontier of selection decisions, which represents different trade-offs across multiple traits. We propose a new approach, which maximizes certain traits while keeping others within desirable ranges. Optimal selection decisions are made using a new version of the look-ahead selection algorithm, which was recently proposed for single trait genomic selection and achieved superior performance with respect to other state-of-the-art selection methods. To demonstrate the effectiveness of the new method a case study is developed using a realistic data set where our method is compared with conventional index selection. Results suggest that the multi-trait look-ahead selection is more effective at balancing multiple traits compared to index selection.

KEYWORDS Multi-trait Genomic Selection; Simulation; Optimization

1. Introduction

Genomic selection (GS), which was initially proposed by Meuwissen *et al.* (2001), is a special form of marker assisted selection (MAS) that simultaneously estimates the effects of genome-wide markers in a training population consisting of genotyped and phenotyped individuals. Selection decisions are based on genomic estimated breeding values (GEBVs) in a breeding population, which are calculated as the sum of the estimated marker effects. The advantages of GS have been demonstrated by simulation and empirical studies (Meuwissen *et al.* 2001; Makowsky *et al.* 2011; Schaeffer 2006; Wang *et al.* 2018; Goddard 2009).

Previous studies have mainly focused on the development of models to improve the accuracy of GEBV prediction. Until recently, few studies have considered alternatives to truncation selection on GEBVs followed by random mating of the selected individuals. These studies have focused on selecting the parents of the next generation by defining new quantitative selection metrics (Goddard 2009; Daetwyler *et al.* 2015; Goiffon *et al.* 2017; Moeinizade *et al.* 2020) or jointly considering selec-

tion and mating decisions (Moeinizade *et al.* 2019; Akdemir and Sánchez 2016). The latter two methods are forms of mate selection (Kinghorn and R.K.Shepherd 1999) that optimize the contributions of potential parents to the next generation based on maximizing a desired breeding objective. Typically, the optimization is performed with respect to the next generation (Kinghorn and Kinghorn 2016; Akdemir and Sánchez 2016). Look-ahead mate selection (LAMS) schemes that optimize parental contributions with respect to grand-progeny (i.e., two generations in the future) have also been proposed in the context of animal breeding (Hayes *et al.* 1998; Shepherd and Kinghorn 1998; Hayes *et al.* 2002).

Moeinizade *et al.* (2019) implemented a LAMS scheme –look-ahead selection (LAS)– in a stochastic simulation framework that seeks to optimize the performance of the best possible progeny in an arbitrarily defined terminal generation. This strategy was shown to outperform conventional genomic selection (Meuwissen *et al.* 2016), optimal haploid value selection (Daetwyler *et al.* 2015), and optimal population value selection (Goiffon *et al.* 2017) using empirical data from a population of maize inbred lines. LAS outperformed previous approaches by achieving more genetic gain and preserving more genetic diversity over the course of a simulated breeding program.

doi: 10.1534/genetics.XXX.XXXXXX

Manuscript compiled: Friday 29th May, 2020

¹Corresponding author's email address: gphu@iastate.edu.

1 Although LAS presents a significant improvement over competing methods, it is confined to single trait genomic selection (ST-GS). Generally, the productivity of a crop variety is dependent on multiple characteristics such as yield, grain quality, and disease resistance. Hence, selection and mating decisions should be based on several different characteristics with potentially different breeding goals. Multi-trait selection poses difficulties for breeders because it often requires balancing competing breeding objectives. Four principle multi-trait genomic selection (MT-GS) strategies have been proposed in the literature: 1) tandem selection, whereby different traits are selected independently in different generations (Burgess and West 1993); 2) independent culling, whereby truncation selection is performed on multiple traits simultaneously with independent thresholds (Hazel 1943); 3) index selection, whereby multiple traits are selected at the same time by constructing an index that is a linear combination of multiple traits (Hazel and Lush 1942; Hazel 1943); and 4) mate selection, whereby multiple traits are selected at the same time by finding Pareto optimal solutions of a mate selection index (Kingham and Kinghorn 2016).

2 Tandem selection, by definition, is not capable of selecting multiple traits simultaneously and is most useful when some traits should be selected in earlier generations than others. Independent culling does perform simultaneous selection but is very sensitive to the truncation points for the different traits. Index selection has become an important method that has been widely used for the development of superior varieties in both animal and plant breeding (Villanueva and Woolliams 1997; Jannink *et al.* 2000; Ivkovich and Koshy 2002; Sharma and Duveiller 2003; Long *et al.* 2006; Yan and Frégeau-Reid 2008). This often takes the form of truncation selection on an index constructed by integrating information on the economic values of the different traits and their phenotypic and additive genetic covariances. Brascamp (1984) provides a concise summary of different selection indices. Mate selection can consider different constraints and breeding goals for multiple traits and evaluates these criteria in the context of a proposed set of matings. Two recent studies in plants have evaluated the use of mate selection on long-term genetic gains (Cowling *et al.* 2019; Suontama *et al.* 2018). Additionally, Akdemir and Sánchez (2016) and Akdemir *et al.* (2019) have developed new mate selection methods for single- and multi-trait scenarios, respectively, with an emphasis on application to plant breeding.

3 An additional challenge in multi-trait selection is the definition of breeding objectives for each trait. For example, a breeder wishing to maximize grain yield might also need to maintain minimum standards for standability and disease resistance and an acceptable range of plant heights. Kempthorne and Nordskog (1959) proposed maintaining a trait at an optimal level by weighting its squared deviations from the optimum. Wilton *et al.* (1968) generalized this approach to include both squares and cross products of multiple traits. Moav and Hill (1966) developed a graphical method to calculate explicitly non-linear indices on two traits. Later, iterative solutions were developed to identify the optimal weights for a non-linear index on an arbitrary number of traits Itoh and Yamada (1988); Pasternak and Weller (1993). However, the general solution for the weights of a non-linear index is dependent on the population mean prior to selection and the intensity of selection (Weller *et al.* 1996). Therefore, the optimum selection index changes each generation and will be different from an index that maximizes gains over multiple generations.

4 In this paper, we propose an extension of the single-trait LAS method to multiple traits with different breeding objectives. The method maximizes a single, main trait while constraining other traits to fall within flexibly defined ranges. It retains the advantages of single-trait LAS derived from considering the impacts of selection, mating, and resource allocation decisions on the performance of individuals in the terminal generation of the breeding program.

2. Materials and Methods

2.1. Data sets

5 A dataset of 5,022 maize recombinant inbred lines (RILs) from the US nested association mapping (US-NAM) (Yu *et al.* 2008) and intermated B73xMo17 (IBM) (Lee *et al.* 2002) populations was used in this study. Best linear unbiased predictors (BLUPs) for total kernel weight were taken from Yang *et al.* (2018). BLUPs for ear height were calculated from the phenotypic data in Kusmec *et al.* (2017) using a mixed model with genotype and environment as random effects. The mixed model was implemented in the R package lme4 (Bates *et al.* 2015).

6 SNPs from Kusmec *et al.* (2017) were thinned using PLINK v1.90b (Chang *et al.* 2015) using the “indep-pairwise” function with a window size of 250 kb, a step size of 50 SNPs, and an LD threshold of 0.6. Thinned SNPs were imputed and phased with Beagle v4.0 (Browning and Browning 2008) using default parameters. This produced 359,826 imputed and phased SNPs. SNP effects for each phenotype were estimated using the BayesB algorithm (Meuwissen *et al.* 2001) implemented in GenSel4 (Fernando and Garrick 2009). Recombination rates were estimated using the genetic map for the US-NAM population (Yu *et al.* 2008) following the procedure outlined in Goiffon *et al.* (2017).

2.2. Simulation design

7 One hundred independent simulations of a ten-generation breeding program were performed using a maize data set. An initial population of 200 individuals was randomly selected from the full data set, and in each generation, 20 individuals were selected to make 10 crosses. More details on the simulation steps are available in Goiffon *et al.* (2017) and Moeinizade *et al.* (2019).

2.3 Single-trait look-ahead selection

8 In this section, we review the look-ahead selection method which was recently proposed for single-trait genomic selection (Moeinizade *et al.* 2019). To make this algorithm more robust, we present an equivalent reformulation of this method and then discuss how this algorithm can be extended for multiple trait settings in the next section.

9 The single-trait look-ahead selection (ST-LAS) method anticipates the consequences of selection and mating decisions over several generations via simulation by quantitatively taking into account recombination frequencies during meiosis. The ST-LAS method has three major contributions to the literature: 1) time management: ST-LAS is the only GS method that takes time constraints into account and is deadline sensitive; 2) mating strategy optimization: the ST-LAS method not only makes the selection decisions but also specifies how to pair the selected individuals for mating; and 3) resource allocation: this method uses a heuristic strategy to allocate more progeny to crosses between more diverse parents to increase the probability of producing high performing individuals.

10 The cornerstone of this method is evaluating a given selection and mating strategy by estimating the distribution of progeny

1 GEBVs in the final generation. By simulating the GEBVs of a
 2 random sample of individuals in the final generation, a breeder
 3 can make better selection and mating decisions. This method can
 4 be formulated as the following optimization model (Moeiniazade
 5 *et al.* 2019):

$$\max_{x,y} f^{\text{LAS}}(x,y,r,\tau) \quad (1)$$

$$\text{s.t.} \quad \sum_{n=1}^N x_n = S \quad (2)$$

$$x_n \in \{0,1\} \quad \forall n \in \{1,\dots,N\} \quad (3)$$

$$x_n = \sum_{j=1}^N y_{n,j} \quad \forall n \in \{1,\dots,N\} \quad (4)$$

$$y_{i,j} \in \{0,1\} \quad \forall i,j \in \{1,\dots,N\} \quad (5)$$

6 This optimization model has two decision variables: x , which
 7 represents the selection strategy, and y , which represents the
 8 mating strategy. Below is a detailed description of the objective
 9 as well as all variables and parameters used in this model:

- 10 • f^{LAS} : The expected GEBV of the best offspring in the termi-
 11 nal generation.
- 12 • x_n : A binary decision variable that shows whether individ-
 13 ual n is selected ($x_n = 1$) or not ($x_n = 0$).
- 14 • $y_{i,j}$: A binary variable that shows whether individual i is
 15 mated with individual j ($y_{i,j} = 1$) or not ($y_{i,j} = 0$).
- 16 • $r \in [0, 0.5]^{L-1}$: The recombination frequency vector.
- 17 • τ : The remaining number of generations ($\tau = T - t$ where
 18 t is the current generation and T is the deadline generation
 19 number).
- 20 • N : The number of individuals in the population.
- 21 • S : The number of individuals that are to be selected out of
 22 the current population.

23 As demonstrated in equation (1), the objective of the ST-LAS
 24 method is dependent on selection (x), mating (y), recombina-
 25 tion frequencies (r), and remaining number of generations (τ).
 26 Constraint (2) states that S individuals are selected from total
 27 N individuals in the population to make $S/2$ crosses (assuming
 28 that S is an even number) and constraint (3) ensures that the
 29 decision variable x , is binary. Constraint (4) ensures that each
 30 selected individual is mated once. Finally, constraint (5) states
 31 that the decision variable y is binary.

32 In this model, evaluation of $f^{\text{LAS}}(x,y,r,T-t)$ is very chal-
 33 lenging because of the uncertainty involved due to recombina-
 34 tion frequencies (r) and also selection (x) and mating (y) deci-
 35 sions over $T-t$ generations. To deal with these challenges, a
 36 simulation optimization algorithm was designed that estimates
 37 and maximizes the LAS objective function by exploring the se-
 38 lection and mating solution space efficiently.

39 2.4 Equivalent formulation of single-trait look-ahead selection

40 According to Moeiniazade *et al.* (2019), the objective of ST-LAS
 41 is to maximize the expected GEBV of the best offspring in the
 42 terminal generation (equation (1)). The *best* offspring can be the
 43 individual with maximum expected GEBV in the final genera-
 44 tion; however, the maximum value does not necessarily repre-
 45 sent the whole distribution. To make the prediction more robust
 46 and reduce the influence of outliers, we present an equivalent re-
 47 formulation of the ST-LAS method (equations (6)-(8)) where the
 48 *best* offspring is defined as the $100\gamma^{\text{th}}$ percentile among predicted
 49 GEBVs of individuals in the terminal generation.

$$\max_{x,y} \quad \phi \quad (6)$$

$$\text{s.t.} \quad \text{Constraints (2), (3), (4), and (5)} \quad (7)$$

$$\Pr[g_1(x,y,r,\tau) \geq \phi] \geq 1 - \gamma \quad (8)$$

Here, ϕ is a threshold value, equivalent to the previous ob-
 50 jective f^{LAS} , which represents the expected GEBV of the best
 51 offspring in the final generation where best is defined as the
 52 $100\gamma^{\text{th}}$ percentile of the simulated GEBV distribution. The new
 53 variables and parameters are defined as follow:
 54

- 55 • ϕ : The expected GEBV of the best offspring in the terminal
 56 generation.
- 57 • $g_1(x,y,r,\tau)$: The expected GEBV of a random progeny in
 58 the terminal generation (for trait 1 which is the only trait in
 59 the case of ST-LAS).
- 60 • γ : A parameter that defines which percentile of the GEBV
 61 distribution is evaluated in the final generation.

62 In this model, constraint (8) states that for a random progeny
 63 in the final generation, the probability of having an expected
 64 GEBV at least equal to the threshold value is greater than or
 65 equal to $1 - \gamma$. For example, for a random sample of 1000
 66 progeny, if $\gamma = 0.98$, then ϕ will evaluate the GEBV of the
 67 top 2% of progeny.

68 2.5 Multi-trait look-ahead selection

69 In this section, we present a new approach for MT-GS prob-
 70 lems to optimize the main goal of a breeding program while
 71 keeping other traits within desired ranges. This new approach,
 72 multi-trait look-ahead selection (MT-LAS), extends the ST-LAS
 73 method to multiple trait settings. It should be noted that the
 74 same resource allocation heuristic from Moeiniazade *et al.* (2019)
 75 is applied to MT-LAS. This resource allocation strategy aims
 76 to preserve more genetic diversity by varying the number of
 77 progeny produced from each cross relative to their breeding
 78 parents genetic diversity.

79 Assume there exists J different traits of which one, $j = 1$ (e.g.,
 80 yield), should be maximized while the other traits, $j \in \{2, 3, \dots, J\}$
 81 (e.g., plant height, ear height, etc.), should satisfy certain criteria.
 82 This problem can be formulated as an optimization model as
 83 follows:

$$\max_{x,y} \quad \phi \quad (9)$$

$$\text{s.t.} \quad \text{Constraints (2)-(5)} \quad (10)$$

$$\Pr[g_1(x,y,r,\tau) \geq \phi | l_j \leq g_j(x,y,r,\tau) \leq u_j, \quad (11)$$

$$\forall j \in \{2, 3, \dots, J\}]$$

$$\geq 1 - \gamma$$

This model shares the same objective and constraints (2),
 84 (3), (4), and (5) with the equivalent reformulation of ST-LAS.
 85 However, constraint (11) is a modification of constraint (8) which
 86 focuses on making sure that traits $j \in \{2, 3, \dots, J\}$ fall into desired
 87 ranges by defining a conditional probability. Below is a detailed
 88 description of all new variables and parameters:
 89

- 90 • $g_j(x,y,r,\tau)$: The expected GEBV of a random progeny in
 91 the terminal generation for trait j where $j \in \{2, 3, \dots, J\}$.
- 92 • l_j : The lower value for trait j .
- 93 • u_j : The upper value for trait j .

This model aims to maximize the expected GEBV of the top 100(1 - γ)% of offspring in the terminal generation for the trait of interest (e.g., yield) among offspring that also meet thresholds with respect to other traits (e.g., plant height, grain quality, etc.). Without loss of generality, $l_j = -\infty$ or $u_j = \infty$ capture the cases when only a lower bound or upper bound should be considered. Note that when only one trait ($j = 1$) is considered, this formulation is equivalent to ST-LAS.

The ST-LAS optimization model was already challenging to solve and after adding a nonlinear and non-convex constraint (11), the computational complexity increases significantly. To overcome this challenge, we redefine constraint (11) by converting the conditional probability on l and u to a penalty that dynamically adjusts the objective function in response to violations of the boundaries. The penalty allows violations of the boundaries that are offset by improvements in the objective function. Take, for example, the case that the decision maker wants to maximize yield while making sure that plant height does not exceed a certain value. What if we could improve yield by slightly violating the height constraint? We want the height constraint to be true, but not at the expense of losing the main objective.

The following mathematical model formulates the problem:

$$\max_{x,y} \quad \phi \quad (12)$$

$$\text{s.t.} \quad \text{Constraints (2)-(5)} \quad (13)$$

$$\theta_j = \Pr[l_j \leq g_j(x, y, r, \tau) \leq u_j], \forall j \in \{2, \dots, J\} \quad (14)$$

$$\Delta = \max(g_j(x, y, r, \tau) - u_j, l_j - g_j(x, y, r, \tau), 0) \quad (15)$$

$$\Pr[h(x, y, r, \tau) \geq \phi] \geq 1 - \gamma \quad (16)$$

$$h(x, y, r, \tau) = \theta_1 g_1(x, y, r, \tau) - \sum_{j=2}^J \frac{(1-\theta_j)}{J-1} \Delta \quad (17)$$

Here, θ_j is the probability that a random progeny is acceptable in the final generation with respect to trait j for $j \in \{2, 3, \dots, J\}$

and $\theta_1 = \frac{\sum_{j=2}^J \theta_j}{J-1}$. The new function $h(x, y, r, \tau)$ is a linear combination of the expected GEBV of a random progeny for trait $j = 1$ and the penalty of violating the desired range for traits $j \in \{2, 3, \dots, J\}$.

Here are some properties of constraints (14)-(17) :

- The term Δ in equation (15) represents the penalty for violating the upper or lower bounds for a random progeny in the terminal generation. As the magnitude of the violation increases, the penalty term increases. In the case of no violation, the penalty becomes 0.
- From equation (17), the term $\sum_{j=2}^J \frac{(1-\theta_j)}{J-1} \Delta$ is the weighted sum of penalties for all traits of $j \in \{2, 3, \dots, J\}$. The weight $(1 - \theta_j)$ is the probability that a random progeny violates the desired range.
- When all the individuals with respect to traits $j \in \{2, 3, \dots, J\}$ (e.g., height) are acceptable, $\theta_1 = 1$ and the focus will be only on the trait of interest (e.g., yield).
- The sum of all weights in equation (17) equals 1 ($\theta_1 + \sum_{j=2}^J \frac{(1-\theta_j)}{J-1} = \frac{\sum_{j=2}^J \theta_j}{J-1} + \sum_{j=2}^J \frac{(1-\theta_j)}{J-1} = 1$).
- The larger that θ_1 is, more weight is placed on the trait of interest (e.g., yield) in selection and mating decisions.

After evaluation of the MT-LAS objective function, the next step is to optimize the model. A similar heuristic algorithm from Moeiniazade *et al.* (2019) is used to optimize the MT-LAS model. This algorithm is defined as follows.

Algorithm 1 Heuristic for optimizing the MT-LAS model

```

1: Select  $S$  random individuals from the population
2: Randomly mate selected individuals
3: Calculate  $\phi$  ( $Vmax \leftarrow \phi$ )
4: Set  $f \in \{1, 2, \dots, S\}$  as list of positions to check
5: Set  $nf \in \{1, 2, \dots, S\}$  as number of positions to check
6: while  $nf \geq 0$  do
7:   Generate  $z \in [1, nf]$  as a random integer
8:    $i \leftarrow$  the  $z^{\text{th}}$  value in  $f$ 
9:    $j \leftarrow$  index of the  $i^{\text{th}}$  individual
10:  Swap  $j$  with every unselected individual from population
11:  Calculate  $\phi_w$  for every possible swap  $w$ 
12:   $VmaxN \leftarrow \max(\phi_w)$ 
13:  if  $VmaxN \leq Vmax$  then
14:    Reject the swap and keep  $j$ 
15:    Remove the  $z^{\text{th}}$  position from  $f$ 
16:  else
17:    Accept the swap
18:     $Vmax = VmaxN$ 
19:     $f \in \{1, 2, \dots, S\} \setminus i$ 
20:     $nf = S - 1$ 
21:  end
22: end

```

2.6 Example with illustration

In this section, we illustrate the MT-LAS method with an example to provide a more intuitive description. Assume that the goal is maximizing yield (trait 1) while ensuring that plant height (trait 2) falls within a desired range. For a given selection and mating strategy at the current generation (t), the look-ahead stochastic simulation predicts the GEBV of individuals in the final generation (T) with respect to both traits as illustrated in Figure 1.

In this example, 20 random progeny are produced in the final generation. The GEBVs of these progeny for both traits are approximated with the look-ahead algorithm. In Figure 1, the green and blue bars represent the GEBVs for each progeny with respect to traits 1 and 2 (i.e., $g_1(x, y, r, \tau)$ and $g_2(x, y, r, \tau)$, respectively). GEBVs for plant height are constrained to fall between 15 and 35. Hence, among all progeny, lines 1, 6, 7, 8, 9, 12, 13, 15, 19, and 20 are not acceptable for plant height. These progeny are distinguished from progeny that meet the height requirements with a cross mark. Because 10 out of 20 individuals meet the height criterion, θ_1 and θ_2 are both 0.5. Finally, the penalty (Δ) and penalized GEBVs for each progeny are calculated using equations (15) and (17), respectively. Penalized GEBVs are plotted as the purple bars in Figure 1.

After sorting the progeny with respect to penalized GEBVs, we can calculate the objective ϕ . Let's assume $\gamma = 0.90$. The 90th percentile among 20 individuals is the third best individual and according to Figure 1, line 14 is the third best individual. Hence we have $\phi = 21$ which is the value of $h(x, y, r, \tau)$ for line 14.

2.7 Data availability

Data are available at Figshare (DOI: 10.25380/iastate.12145752).




Figure 1 The look-ahead simulation illustration for MT-LAS method. In this example, the population consists of 16 individuals. In generation t , 8 individuals are selected from the population and mated accordingly to make 4 crosses. Each breeding parent produces one progeny in generation $t+1$ and from generation $t+1$ to $T-1$ all progeny are crossed with each other in the same generation, each producing one progeny. Then, the look-ahead objective can be approximated by taking a random sample of progeny in generation T . In this example, 20 lines are produced and the GEBV of each individual with respect to traits 1 and 2 are measured and visualized with green and blue bars respectively. Our goal is to maximize trait 1 after $T-t$ generations while making sure that trait 2 does not exceed a certain value of $u = 35$ and is not less than $l = 15$. We observe that 10 individuals among 20 are not acceptable. The progeny with acceptable values for bounded trait are distinguished with check marks. The penalized GEBVs are calculated and represented as purple bars and calculation of the objective ϕ is demonstrated for a given γ .

3. Results

In this section, we present a case study with real data. Without loss of generality, two traits—total kernel weight (TKW) and ear height (EHT)—are used in this case study. Our objective is to maximize TKW given a constraint on EHT.

We first present the performance of ST-LAS where the goal is to maximize TKW only. In this way, we can observe the behavior of EHT versus TKW in the absence of any constraints on EHT. Then, we investigate truncation selection on a selection index for TKW and EHT with different choices of weights. However, this does not allow keeping a trait within a specified range. Hence, we define a penalized index by assigning a negative weight on the absolute deviations from the specified range. The penalized index is used as a benchmark against the performance of MT-LAS. Finally, we present the MT-LAS results and compare the effectiveness of MT-LAS to that of the penalized index.

Single-trait look-ahead selection - Maximizing TKW

In this section, we investigate the behavior of EHT over ten generations when the objective is to maximize TKW and there is no constraint on EHT. This will help provide reasonable bounds for EHT when testing the MT-LAS algorithm.

Figure 2 (A) presents the total kernel weight and ear height GEBVs over ten generations for a single simulation when selection is only on TKW. Over ten generations the mean GEBV of TKW increases from 2.78 to 40.25 with a maximum of 47.09. For EHT, the range of GEBVs changes from $[-21.87, 25.42]$ to $[-1.68, 14.77]$. Figure 2 (B) presents the minimum, mean, and maximum GEBVs of both traits over ten generations averaged over 100 simulation replicates. On average, the GEBVs of EHT fall in a range of $[-2.73, 16.54]$ in the final generation.

Index selection - Maximizing TKW and EHT with assigned weights

A selection index is a linear combination of traits according to some weighting scheme. Typically, truncation selection is applied to the index. Here, we construct an index for TKW and EHT where the index is the weighted sum of the GEBVs for each trait ($W_{TKW}GEBV_{TKW} + W_{EHT}GEBV_{EHT}$) and truncation selection is applied to the index. Let W_{TKW} and W_{EHT} be the weights placed on the GEBVs for total kernel weight and ear height, respectively. Weights are chosen from the real numbers between -1 and 1. It should be noted that, in this scheme, we are selecting for larger values of both TKW and EHT. Placing a negative weight on a trait selects for smaller values and produces progress in the opposite direction to that under strictly positive weights. Figure 3 presents the average GEBVs over ten generations averaged over 100 replicate simulations under index selection with varying weights, including the case where the weight on EHT is negative.

As expected, increased weight for EHT (positive or negative) negatively impacts the efficiency of selection for TKW. The mean GEBVs for both traits change in the direction of their assigned weights over time, indicating the lack of strong genetic correlations between TKW and EHT. The highest mean GEBV for TKW (34.67) is achieved by selection solely on TKW ($W_{TKW} = 1, W_{EHT} = 0$). Table 1 provides the minimum, mean, and maximum GEBVs for TKW in the final generation under the different choices for weights. It should be noted that the maximum GEBV for TKW achieved after 10 generations of selection is less than that achieved using ST-LAS (36.05 vs. 44.51). This

considerable impact on response is due to the fact that the look-ahead selection focuses on maximizing the expected GEBV of the best offspring in the terminal generation, considering uncertainty in recombination in each generation whereas truncation selection on GEBVs focuses on maximizing the genetic gain in the next generation. Additionally, look-ahead selection selects pairs of individuals as a group and recognizes the importance of mating.

Table 1 Summary statistics of population GEBV values in generation 10 averaged over 100 replicate simulations for TKW using conventional genomic selection with different weights (index selection). These results are based on simulations in Figure 3.

W_{TKW}	W_{EHT}	Min	Mean	Max
0	± 1	0.22	2.81	5.41
0.1	± 0.9	1.86	4.63	7.53
0.2	± 0.8	4.23	7.08	9.86
0.3	± 0.7	7.21	10.28	13.3
0.4	± 0.6	12.07	15.30	18.43
0.5	± 0.5	18.53	21.68	24.69
0.6	± 0.4	25.86	28.77	31.6
0.7	± 0.3	30.21	32.57	34.85
0.8	± 0.2	32.00	33.81	35.49
0.9	± 0.1	32.69	34.11	35.48
1	0	33.29	34.67	36.05

Penalized index selection - Maximizing TKW with a constraint on EHT

In this section, we reformulate the index selection to be able to specify a desired range for the secondary trait. This enables a direct comparison to the MT-LAS method. After applying ST-LAS to TKW, the GEBVs for EHT in the final generation fell between -2.73 and 16.54 . We subsequently investigated three cases where EHT is constrained to fall outside this range of variation. The three cases are as follows:

- Case 1: $l = 20, u = 30$
- Case 2: $l = -15, u = -5$
- Case 3: $l = 45, u = +\infty$

Similar to the use of a quadratic index to approach an optimum phenotype (Kempthorne and Nordskog 1959; Wilton et al. 1968), we define an index that penalizes the absolute deviations from a desired range. The constructed index is formulated as $W_{TKW}GEBV_{TKW} - W_{EHT} \max(l - GEBV_{EHT}, 0, GEBV_{EHT} - u)$. Weights are chosen from the real numbers between 0 and 1, constrained to sum to unity. Figure 4 presents the average GEBVs over ten generations averaged over 100 replicate simulations under penalized index selection for three different cases. These results are compared against the index selection without penalization from Figure 3. We observe that the non-penalized index selection cannot satisfy the ear height criterion. As expected, over multiple generations of selection the GEBV of TKW increases and the penalty term accommodates keeping EHT within




Figure 2 (A) Population GEBVs of EHT versus TKW for one simulation replicate over ten generations when selection and mating decisions are optimized using ST-LAS algorithm with an objective of maximizing TKW. Each generation includes 200 individuals represented by stars and different colors are distinguishing between generations. The final generation has a minimum, mean, and maximum of 34.36, 40.25, 47.09 for TKW and $-1.68, 7.17, 14.77$ for EHT respectively. (B) Minimum, mean and maximum GEBVs of TKW and EHT over ten generations averaged over 100 simulation replicates. Selection and mating decisions are optimized using ST-LAS algorithm with an objective of maximizing TKW. The final generation has a minimum, mean, and maximum of 33.30, 39.04, 44.51 for TKW and $-2.73, 7.00, 16.54$ for EHT respectively.




Figure 3 Index selection considering different weights for TKW and EHT averaged over 100 simulation replicates. The mean GEBV of individuals over ten generation are calculated given a pair of weights for two traits. The absolute values of the weights add up to 1. Each curve demonstrates the mean GEBV of individuals (represented by markers) over ten generations for assigned weights.

1 the specified range. For case 1 and case 2, the EHT criterion is
2 satisfied when $W_{\text{EHT}} \geq 0.3$. However, for case 3, the criterion
3 cannot be satisfied even with the penalized index selection be-
4 cause the bound represents an extreme case. The behavior of
5 case 3 is very similar to the index selection without penalization.

6 **Multi-trait look-ahead selection - Maximizing TKW with a con-** 7 **straint on EHT**

8 The MT-LAS method aims to maximize genetic gain in a tar-
9 get trait while ensuring that one or more secondary traits fall
10 within specified boundaries. Here, we maximize TKW subject
11 to constraints on EHT for three different cases.

12 Population GEBVs over ten generations for one simulation
13 replicate are presented in Figure 5 (A) and the average of 100
14 simulation replicates are presented in Figure 5 (B). For cases 1
15 and 2, the GEBVs for EHT of $\sim 90\%$ of the individuals in the
16 final generation fall within the specified boundaries. For case 3,
17 only a lower bound on EHT GEBV was specified. This bound
18 represents an extreme case where index selection is unable to
19 achieve the lower bound even when selecting solely on EHT.
20 However, MT-LAS is able to exceed the bound with $\sim 50\%$ of
21 the population falling into the acceptable range.

22 **Comparison - Performance of MT-LAS against penalized index** 23 **selection**

24 Figure 6 compares the performance of MT-LAS with the results
25 of non-penalized and penalized index selection using weights
26 that produced results satisfying the desired ranges for 3 cases.
27 We also show that ST-LAS for TKW exceeds the performance of
28 truncation selection on TKW alone ($W_{\text{TKW}} = 1, W_{\text{EHT}} = 0$). For
29 both cases 1 and 2, MT-LAS is able to produce populations that
30 surpass the performance of the comparable index selection scenar-
31 ios with respect to TKW and also keep almost all individuals
32 within the specified boundaries for EHT. For case 3, the highest
33 EHT achieved by index selection with or without penalization
34 cannot satisfy the desired range criterion. However, MT-LAS
35 not only achieves the expected EHT, but also improves the TKW
36 considerably.

37 Overall, using MT-LAS with optimization of selection and
38 mating decisions and a soft penalty on ear height improves the
39 response. It should be noted that the distributions of look-ahead
40 methods are quite different from index selection. As shown in
41 Figure 6 the look-ahead methods achieve wider distributions in
42 the terminal generation.

43 Figures 7 and 8 display the standard deviation of population
44 GEBVs over ten generations for 100 simulation replicates and
45 compare the performance of MT-LAS/ST-LAS with index selec-
46 tion. As expected, look-ahead methods maintain more genetic
47 variance than index selection indicating that there is greater
48 room for population improvement after 10 generations. Further-
49 more, the genetic correlations between two traits are presented
50 over time for one simulation replicate which indicate the lack
51 of strong correlation between TKW and EHT (see Figure 9 in
52 Appendix).

53 **Discussion**

54 The production of a crop variety depends on multiple character-
55 istics such as grain quality, yield, and drought resistance which
56 are subject to different breeding objectives. In this study, we
57 proposed a new multi-trait selection approach using genomic
58 information that maximizes genetic gain with respect to a focal
59 trait while controlling the variation in multiple secondary traits.

60 To demonstrate the effectiveness of the proposed method, we
61 conducted a case study using real data where MT-LAS is com-
62 pared with index selection with varying weights. In this case
63 study, the goal was to maximize total kernel weight while con-
64 straining ear height. Three different cases with varying bounds
65 were investigated, and the results suggested that MT-LAS was
66 more effective at balancing multiple traits than index selection.

67 Fundamentally, the MT-LAS algorithm surpassed conven-
68 tional index selection because of four reasons. The first reason
69 is the **satisfiability** of this method. MT-LAS automatically and
70 dynamically balances multiple traits and is able to optimize
71 selection and mating decisions in a way that satisfies the con-
72 straints for bounded traits while simultaneously maximizing
73 the main trait of interest in the terminal generation. For two of
74 our three scenarios, the penalized index was able to satisfy the
75 constraints on the bounded trait but at the cost of reduced per-
76 formance in the maximized trait. Moreover, with index selection,
77 it may not be possible to achieve some values for the bounded
78 traits without mate selection. For example, in case 3, we inves-
79 tigate the performance of MT-LAS with a lower bound of 45
80 which is not reached with either non-penalized or penalized
81 index selection (Figures 3 and 4).

82 The second advantage of MT-LAS is its **dynamic adjustabil-**
83 **ity**. The MT-LAS method places more emphasis on feasibility re-
84 quirements (having individuals that meet the thresholds for the
85 bounded traits) when most of the individuals are not predicted
86 to fall within the bounds for the bounded traits in the terminal
87 generation. On the other hand, this algorithm focuses on the
88 main trait when most of the individuals become acceptable for
89 the bounded trait. Overall, selection and mating decisions are
90 dynamically adjusted in every generation by making a trade
91 off between optimizing the main goal and reaching the desired
92 range for the bounded traits.

93 A third benefit of MT-LAS is its **interpretability**. By defining
94 the weights in terms of bounds on the desired values of the
95 trait, MT-LAS provides an intuitive description of the breeding
96 objective on the original measurement scale.

97 A fourth benefit of MT-LAS is its **time-awareness**. As op-
98 posed to classical index selection, which maximizes genetic merit
99 in the next generation, MT-LAS maximizes genetic merit in an
100 arbitrary terminal generation. This is similar to work on look-
101 ahead mate selection in animal breeding (Hayes *et al.* 1998; Shep-
102 herd and Kinghorn 1998; Hayes *et al.* 2002) where the quantity to
103 be maximized is the genetic merit of grand-progeny. Addition-
104 ally, this shift alleviates the difficulties posed by the dependence
105 of classical non-linear indices on the current generation mean
106 and intensity of selection which can cause such an index to be
107 non-optimal over multiple generations (Weller *et al.* 1996).

108 The main contribution of MT-LAS is constraints (14), (16),
109 and (17), which allow the algorithm to dynamically adjust the
110 objective function according to the progress of the current popu-
111 lation. Future research is needed to more fully characterize
112 the MT-LAS algorithm and address the limitations of this study.
113 First, the current paper only considers two traits, although the
114 model is formulated for J traits. Further simulations to explore
115 the behavior of the algorithm when constraining >1 trait are
116 desirable. Second, the hyper-parameter γ plays a crucial role
117 in identifying the optimal selection and mating decisions. In
118 this study we selected γ after experimenting with several values.
119 Future work is needed to design systematic methods for opti-
120 mizing this parameter. Third, the objective of the look-ahead
121 selection relates to the final generation and future research can




Figure 4 Penalized index selection considering different weights for TKW and EHT averaged over 100 simulation replicates for three different cases. Each curve demonstrates the mean GEBV of individuals (represented by markers) over ten generations for assigned weights. The transparent curves in the background present the index selection results without penalization and the red dashed lines are the decision boundaries.

1 focus on designing new selection methods that also consider
 2 intermediate generations in the objective. Finally, we based our
 3 simulations on a single data set from a single crop organism.
 4 Further simulations considering more diverse populations are
 5 necessary to demonstrate the general applicability of MT-LAS.

6 Acknowledgements

7 This work is supported by Agriculture and Food Research Initia-
 8 tive Grant no. 2017-67007-26175/Accession No. 1011702 from
 9 the USDA National Institute of Food and Agriculture. Any opin-
 10 ions, findings, conclusions, or recommendations expressed in
 11 this publication are those of the author(s) and do not necessarily
 12 reflect the view of the U.S. Department of Agriculture. This
 13 work is also supported by the Plant Sciences Institute’s Faculty
 14 Scholars program at Iowa State University.

15 Literature Cited

16 Akdemir, D., W. Beavis, R. Fritsche-Neto, A. K. Singh, and
 17 J. Isidro-Sánchez, 2019 Multi-objective optimized genomic
 18 breeding strategies for sustainable food improvement. *Heredity*
 19 **122**: 672.
 20 Akdemir, D. and J. I. Sánchez, 2016 Efficient breeding by genomic
 21 mating. *Frontiers in Genetics* **7**: 210.
 22 Bates, D., M. Mächler, B. Bolker, and S. Walker, 2015 Fitting
 23 linear mixed-effects models using lme4. *Journal of Statistical*
 24 *Software*, **67**(1): 1-48. doi: 10.18637/jss.v067.i01 .
 25 Brascamp, E., 1984 Selection indices with constraints. *Anim.*
 26 *Breed. Abst.* **52**: 645–654.
 27 Burgess, J. C. and D. West, 1993 Selection for grain yield follow-
 28 ing selection for ear height in maize. *Crop Science* **33**: 679–682.
 29 Chang, C. C., C. C. Chow, L. C. Tellier, S. Vattikuti, S. M. Purcell,
 30 *et al.*, 2015 Second-generation plink: rising to the challenge of
 31 larger and richer datasets. *Gigascience* **4**: 7.
 32 Cowling, W. A., L. Li, K. H. Siddique, R. G. Banks, and B. P.
 33 Kinghorn, 2019 Modeling crop breeding for global food se-

curity during climate change. *Food and Energy Security* **8**:
 e00157.
 35
 36 Daetwyler, H. D., M. J. Hayden, G. C. Spangenberg, and B. J.
 37 Hayes, 2015 Selection on optimal haploid value increases ge-
 38 netic gain and preserves more genetic diversity relative to
 39 genomic selection. *Genetics* **200**: 1341–1348.
 40 Fernando, R. and D. Garrick, 2009 Gensel—user manual for
 41 a portfolio of genomic selection related analyses. Technical
 42 report. Available at: <http://biggs.ansci.iastate.edu/bigsgui>. Ac-
 43 cessed: June 13, 2017. .
 44 Goddard, M., 2009 Genomic selection: prediction of accuracy
 45 and maximisation of long term response. *Genetica* **136**: 245–
 46 257.
 47 Goiffon, M., A. Kusmec, L. Wang, G. Hu, and P. S. Schnable, 2017
 48 Improving response in genomic selection with a population-
 49 based selection strategy: Optimal population value selection.
 50 *Genetics* **206**: 1675–1682.
 51 Hayes, B., R. Shepherd, and S. Newman, 2002 Look ahead mate
 52 selection schemes for multi-breed beef populations. *Animal*
 53 *Science* **74**: 13–23.
 54 Hayes, B., R. Shepherd, S. Newman, and B. Kinghorn, 1998 A
 55 tactical approach to improving long term response in across
 56 breed mating plans .
 57 Hazel, L. and J. L. Lush, 1942 The efficiency of three methods of
 58 selection. *Journal of Heredity* **33**: 393–399.
 59 Hazel, L. N., 1943 The genetic basis for constructing selection
 60 indexes. *Genetics* **28**: 476–490.
 61 Itoh, Y. and Y. Yamada, 1988 Linear selection indices for non-
 62 linear profit functions. *Theoretical and Applied Genetics* **75**:
 63 553–560.
 64 Ivkovich, M. and M. Koshy, 2002 Optimization of multiple trait
 65 selection in western hemlock (*tsuga heterophylla* (raf.) sarg.)
 66 including pulp and paper properties. *Annals of Forest Science*
 67 **59**: 577–582.
 68 Jannink, J.-L., J. Orf, N. R. Jordan, and R. G. Shaw, 2000 Index se-
 69 lection for weed suppressive ability in soybean. *Crop Science*
 70 **40**: 1087–1094.




Figure 5 (A) GEBVs of individuals over ten generations for one simulation replicate. Optimal selection and mating decisions were made using the MT-LAS method in all three cases. Generations are distinguished with different colors. Over multiple generations of selection, the GEBV of TKW increases and the GEBV of EHT falls within the desired range. The red dashed lines are the decision boundaries and the arrows demonstrate the direction for which the condition is satisfied. (B) Minimum, mean and maximum GEBVs over ten generations averaged over 100 simulation replicates. The blue markers in the middle of cross marks are the mean GEBVs and the end of the cross marks represent minimum and maximum GEBVs.




Figure 6 Comparison of MT-LAS, ST-LAS and Index selection methods. The mean GEBVs of population over ten generations are averaged over 100 simulation replicates and represented for two traits. Furthermore, the minimum and maximum GEBVs in the final generation are demonstrated using the cross marks. The green bar specifies the boundaries.




Figure 7 Standard deviations of total kernel weight GEBVs over time averaged for 100 simulation replicates.




Figure 8 Standard deviations of ear height GEBVs over time averaged for 100 simulation replicates.

- 1 Kempthorne, O. and A. W. Nordskog, 1959 Restricted selection
2 indices. *Biometrics* **15**: 10–19.
- 3 Kinghorn, B. and A. Kinghorn, 2016 Instructions for matesel.
- 4 Kinghorn, B. and R.K.Shepherd, 1999 Mate selection for the
5 tactical implementation of breeding programs. *Proceedings*
6 *of the Advancement of Animal Breeding and Genetics* **13**:
7 130–133.
- 8 Kusmec, A., S. Srinivasan, D. Nettleton, and P. S. Schnable, 2017
9 Distinct genetic architectures for phenotype means and plas-
10 ticities in *zea mays*. *Nature Plants* **3**: 715.
- 11 Lee, M., N. Sharopova, W. D. Beavis, D. Grant, M. Katt, *et al.*,
12 2002 Expanding the genetic map of maize with the intermated
13 b73× mo17 (ibm) population. *Plant Molecular Biology* **48**:
14 453–461.
- 15 Long, J., J. B. Holland, G. P. Munkvold, and J.-L. Jannink, 2006
16 Responses to selection for partial resistance to crown rust in
17 oat .
- 18 Makowsky, R., N. M. Pajewski, Y. C. Klimentidis, A. I. Vazquez,
19 C. W. Duarte, *et al.*, 2011 Beyond missing heritability: predic-
20 tion of complex traits. *PLoS Genetics* **7**: e1002051.
- 21 Meuwissen, T., B. Hayes, and M. Goddard, 2016 Genomic selec-
22 tion: A paradigm shift in animal breeding. *Animal Frontiers*
23 **6**: 6–14.
- 24 Meuwissen, T. H. E., B. J. Hayes, and M. E. Goddard, 2001 Predic-
25 tion of total genetic value using genome-wide dense marker
26 maps. *Genetics* **157**: 1819–1829.
- 27 Moav, R. and W. Hill, 1966 Specialised sire and dam lines. iv.
28 selection within lines. *Animal Science* **8**: 375–390.
- 29 Moeinizade, S., G. Hu, L. Wang, and P. S. Schnable, 2019 Op-
30 timizing selection and mating in genomic selection with a
31 look-ahead approach: An operations research framework. *G3*:
32 *Genes, Genomes, Genetics* pp. g3–200842.
- 33 Moeinizade, S., M. Wellner, G. Hu, and L. Wang, 2020
34 Complementarity-based selection strategy for genomic selec-
35 tion. *Crop Science* **60**: 149–156.
- 36 Pasternak, H. and J. Weller, 1993 Optimum linear indices for
37 non-linear profit functions. *Animal Science* **56**: 43–50.
- 38 Schaeffer, L., 2006 Strategy for applying genome-wide selection
39 in dairy cattle. *Journal of Animal Breeding and Genetics* **123**:
40 218–223.
- 41 Sharma, R. and E. Duveiller, 2003 Selection index for improving
42 *helminthosporium* leaf blight resistance, maturity, and kernel
43 weight in spring wheat. *Crop Science* **43**: 2031–2036.
- 44 Shepherd, R. and B. Kinghorn, 1998 A tactical approach to the de-
45 sign of crossbreeding programs. In *Proceedings of the sixth world*
46 *congress on genetics applied to livestock production*, volume 25,
47 pp. 431–438.
- 48 Suontama, M., B. Kinghorn, W. Cowling, and H. Dungey, 2018
49 Tactical desired gains for control of red needle cast in radiata
50 pine under optimal contributions selection .
- 51 Villanueva, B. and J. Woolliams, 1997 Optimization of breeding
52 programmes under index selection and constrained inbreed-
53 ing. *Genetics Research* **69**: 145–158.
- 54 Wang, L., G. Zhu, W. Johnson, and M. Kher, 2018 Three new
55 approaches to genomic selection. *Plant Breeding* **137**: 673–681.
- 56 Weller, J., H. Pasternak, and A. Groen, 1996 Selection indices for
57 non-linear breeding objectives, selection for optima. *Interbull*
58 *Bulletin* .
- 59 Wilton, J., D. A. Evans, and L. Van Vleck, 1968 Selection indices
60 for quadratic models of total merit. *Biometrics* pp. 937–949.
- 61 Yan, W. and J. Fréreau-Reid, 2008 Breeding line selection based
62 on multiple traits. *Crop Science* **48**: 417–423.
- Yang, J., R. K. Ramamurthy, X. Qi, R. L. Fernando, J. C. Dekkers, 63
et al., 2018 Empirical comparisons of different statistical mod- 64
els to identify and validate kernel row number-associated 65
variants from structured multi-parent mapping populations 66
of maize. *G3: Genes, Genomes, Genetics* **8**: 3567–3575. 67
- Yu, J., J. B. Holland, M. D. McMullen, and E. S. Buckler, 2008 68
Genetic design and statistical power of nested association 69
mapping in maize. *Genetics* **178**: 539–551. 70

1 **Appendix**

2 **Genetic correlations**

3 Figure 9 demonstrates population GEVVs for one simulation replicate over ten generations for different methods. The standard
 4 deviation of population GEVVs for TKW and EHT are presented over time. It is observed that almost in every generation the
 5 population has higher genetic variation for both traits when selection and mating decisions are optimized using look-ahead methods.
 6 Furthermore, the genetic correlations between two traits are presented over time which shows these two traits are correlated with a
 7 low degree.



Figure 9 Comparison of the population performance for MT-LAS, ST-LAS and index selection methods over ten generations for one simulation replicate. The gray bars specify boundaries. Each box has three numbers including standard deviation of population GEVVs for trait 1 and trait 2 as well as the correlation between two traits from top to bottom respectively.

8 **Repeatability of the results**

9 The simulations are stochastic because they model stochastic recombination events. Figure 10 (Right) depicts the distribution of
 10 breeding values in the final generation for 100 simulations using the same starting population but different random seeds. The left
 11 panels provide a closer look at the first 10 simulations. As expected, there is variation around the average performance across all
 12 simulations. The average of the first 10 simulations is similar to the average of all 100 simulations, suggesting that the results are
 13 repeatable.




Figure 10 Population GEV box-plots for 10 and 100 independent simulations ((Left) and (Right) respectively). Selection and mating decisions are optimized using MT-LAS method (with an objective of maximizing TKW and having a constraint on EHT (lower-bound 20 and upper-bound 30, similar to case 1)). The purple dashed line demonstrates the average of GEVs across all simulations.