

Exploring harm in psychotherapy: Perspectives of clinicians working with children and young people

Barbara Castro Batic¹ | Dan Hayes^{1,2} 

¹University College London, London, UK

²Anna Freud Centre, London, UK

Correspondence

Dan Hayes, Anna Freud Centre, London, UK.
Email: Daniel.Hayes@annafreud.org

Abstract

Aims: The potential for harm to occur from talking therapies has been acknowledged in academic literature. However, there is a paucity of research when it comes to exploring this phenomenon when working with young patients. This study explores clinicians' perspectives on harm from talking therapies when working with children and young people.

Method: Eleven clinicians were interviewed on the types of harm that could occur from talking therapies, as well as the potential mechanisms. Data were analysed inductively using thematic analysis.

Results: Two themes were identified around types of harm: 'clinical deterioration' and 'retraumatisation'. Additionally, four groups of mechanisms were identified: 'Administrative factors', 'Relationship factors', 'Therapist factors' and 'Contextual factors'.

Discussion: Clinicians are able to identify some specific types of harm when working with children and young people and understand how these could occur. The clinical implications of these findings are explored, along with limitations and directions for future research.

KEYWORDS

adolescent, adverse effects, children, harm, psychotherapy, talking therapies

1 | INTRODUCTION

1.1 | Psychotherapy: What could go wrong?

The potential for harm to occur from therapy has been recognised for decades (Barlow, 2010). Yet despite this recognition, the field has not progressed sufficiently (Parry, Crawford, & Duggan, 2016). One reason is the lack of consensus around what is defined as harm and how it links with other frequently used terms, such as adverse events, unwanted effects, negative effects, side effects and

clinical deterioration (Klatte, Strauss, Flückiger, & Rosendahl, 2018; Linden, 2013; Parry et al., 2016; Vaughan, Goldstein, Alikakos, Cohen, & Serby, 2014).

Despite this, how harm can be caused has been previously explored in adult patients (Dimidjian & Hollon, 2010; Duggan, Parry, McMurrin, Davidson, & Dennis, 2014; Linden, 2013). The causes are likely to be multifactorial and to happen because of a combination of patient, clinician and therapeutic factors (Hardy et al., 2019; Jonsson, Johanson, Nilsson, & Lindblad, 2016; Mohr, 1995). For example, therapist factors interact with patient

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Counselling and Psychotherapy Research* published by John Wiley & Sons Ltd on behalf of British Association for Counselling and Psychotherapy

factors, which, in turn, interact with variables from the therapy process (Mohr, 1995).

1.2 | Models to understand or classify harm

Early attempts to classify harm from talking therapies stem from Lilienfeld's (2007) concept of Potentially Harmful Treatments (PHTs). Treatments were classified as PHTs if they met three criteria: (a) demonstrated psychological or physical harm to clients or others, (b) had enduring harmful effects, and (c) harmful effects were replicated by independent research groups. However, a limitation of this approach is that it focuses on classifying treatments as a whole as potentially harmful. This leaves out the iatrogenic effects which could be experienced in a treatment by an individual patient (Blease, Lilienfeld, & Kelley, 2016; Hardy et al., 2019).

Subsequently, Linden (2013) developed a model for classifying side effects from therapy. These started off with unwanted effects (UEs), which were defined as any negative event that occurred during therapy. If the event was specifically related to treatment, then this was an adverse treatment reaction (ATR), whilst if a treatment was applied or given incorrectly, this referred to a malpractice reaction (MPR). Clinical deterioration was kept separate, as it may or may not be related to the other types of harm outlined by Linden (2013). Along with this classification, a checklist was developed to help clinicians identify possible unwanted events of psychotherapy (UE-ATR Checklist; Linden, 2013). It considers the context where the unwanted event emerged (e.g. diagnostic procedures), its severity, the extent to which it is related to treatment, and the area affected (e.g. work, family relationships). Whilst useful, it does not include factors which may be relevant for young people, such as school or college.

More recently, a model has been developed focusing on risk factors for negative experiences during psychotherapy (Hardy et al., 2019). Drawing on empirical data, the authors proposed that a 'lack of fit' between patients, therapists and service structures could lead to tensions in 'Safety and containment', as well as 'Power and Control'. This could cause strain and poor engagement and, for some patients, result in feelings of regret, not being able to cope, hopelessness, lacking confidence, and feeling like a failure. This model, in contrast with previous ones, explicitly considers the patient's experience and complements it with the clinician's. However, the exact relationships between these causal links and how they operate to cause negative experiences needs further validation.

The above models, whilst useful in aiding understanding of how therapy can be harmful or negative, are more tailored for use in adult populations. For example, the last model (Hardy et al., 2019) was based on adult patient data. This does not take into account the unique considerations which children and young people face, which include both physical (e.g. developmental capacity) and psychological vulnerabilities (e.g. power imbalances, multiple stakeholders and mental capacity) which may result in different as well as more severe types of harm (Mercer, 2017).

1.3 | Literature around harm from psychological therapies in adult populations

Literature around harm has often focused on the prevalence of adverse or unwanted events (Moritz et al., 2015, 2018; Schermuly-Haupt, Linden, & Rush, 2018). Recent estimates in patients with depression found that 39% reported at least one side effect (Moritz et al., 2018), whilst for patients with OCD, 93% of patients reported one side effect (Moritz et al., 2015). In another study focusing on unwanted and side effects from CBT, negative effects on well-being/increased distress as a result of treatment was reported in 27% of patients, a worsening in symptoms in 9% of patients, and strains in family relationships in 6% of participants (Schermuly-Haupt et al., 2018). Moreover, 21% of patients reported these negative outcomes as severe or very severe, whilst a further 5% reported these as persistent.

Clinicians' perspectives of negative effects have also been explored. In a study with 74 licensed practitioners, different types of negative effects were identified, and later classified as 'short term negative effects', 'no treatment effect', 'deterioration', 'dependency', and 'impact on other life domains' (Bystedt, Rozental, Andersson, Boettcher, & Carlbring, 2014). Causal factors for harm were also identified. These included 'incompetence and inadequately applied methods', 'failed ethical judgement and professional conduct', and 'discontinuing treatment'. However, this study is limited to the results of clinicians who practiced CBT, and how this translates to other talking treatments remains unclear.

Bowie, McLeod, and McLeod (2016) carried out interviews with therapists, who as clients experienced unhelpful therapy. Four superordinate themes were gathered: 'sourcing a therapist', 'the therapist working to their own agenda', 'the pivotal moment—deciding not to continue' and 'the impact of receiving unhelpful therapy'. The authors concluded that unhelpful therapy may be related to the breakdown of the therapeutic relationship and that there was a difficulty in letting the therapist know about their unhelpful experience. Nevertheless, the study only included participants that had completed their own psychotherapy training and had private therapy, which would differ in terms of the power dynamic and stakeholders involved when it comes to therapy with children and young people.

1.4 | Literature around harm from psychological therapies in children and young people

There is a paucity of studies when it comes to harm from talking therapies with children and young people. One review explored how therapies could be classified as harmful by using the concept of PHT combined with four specific adverse childhood experiences: physical hurt or humiliation, physical abuse, not feeling loved or important, and neglect (Mercer, 2017). Using this framework, four therapies were identified as harmful: Conditioning/Operant Punishment, Holding Therapy/Attachment Therapy and Diagnosis, Festhaltenstherapie (Holding Time, Prolonged Parent-Child Embrace)

and Conversion therapy (Mercer, 2017). The author posited whether the identified therapies should be made less readily available or even prohibited. Whilst useful, this study does not take into account how harm or side effects may occur for children and young people in more mainstream therapies.

To date, only one qualitative study on this topic has been conducted (Jonsson et al., 2016). This study explored Swedish practitioners' experiences of possible adverse effects of psychological therapy with children and adolescents. Four themes were classified from the interviews: 'Vagueness of the concept', 'Psychotherapist-client interaction', 'Consequences of the child' and 'Family effects'. The authors concluded that the concept of harm appeared unfamiliar and vague to many practitioners, and as a result, these may go unnoticed (Jonsson et al., 2016). Whilst helpful in beginning to aid understanding of how therapy might be harmful, this study was skewed towards therapists who mainly practiced within a CBT framework. Thus, how practitioners of other therapeutic domains view harm is unexplored.

Given the paucity of research in the field and the unique considerations which children and young people face when coming to therapy, further research is needed, both with other types of clinicians and within a UK context. As such, the aim of this study is to explore UK clinicians' perspectives on harm when working with children and young people. Questions will focus both on the types of harm that can occur as well as how it occurs (e.g. mechanisms).

2 | METHOD

2.1 | Participants

To be eligible to participate in the study, participants had to meet the following criteria:

- a. Be currently practicing a therapeutic approach in the UK
- b. Work with children and/or adolescents (up to 25 years old)
- c. Have a good grasp of English
- d. Be willing to take part in an interview, either face-to-face or over the phone.

Overall, 11 clinicians were interviewed. This included four males and seven females. Nine clinicians outlined that they worked within an integrative framework, one within a CBT framework, and one within a systemic framework. Clinicians had been practising for a 17.5 years on average ($SD = 9.42$).

2.2 | Procedure

A semi-structured interview schedule was constructed to explore clinicians' notions of harm. Unlike previous research (Jonsson et al., 2016), a definition of harm was not provided at the start of the interview in order to minimise bias and understand how clinicians'

thought of harm and how it could occur (see Appendix A for the developed interview schedule).

Participants were recruited by convenience sampling via social media and through The Anna Freud National Centre for Children and Families networks/newsletters. Clinicians who expressed an interest in taking part were provided with an information sheet and a time was arranged to follow up to see if they had any questions and wanted to participate. Overall, one clinician did not proceed with the study and did not provide a reason why. Those that agreed to take part signed a consent form which was either given to the researcher in person (for face-to-face interviews) or via post/email (if the interview was conducted on the phone). All interviews were undertaken by the primary author (BCB) as part of their MSc dissertation and audio recorded. After taking place, interviews were transcribed verbatim.

2.3 | Data analysis

Interviews were analysed with an inductive thematic analysis (Braun & Clarke, 2006) using NVivo 12 (QSR International, 2018). Analysis was undertaken in a series of steps, which included familiarising oneself with the data, the generation of codes, searching for themes, the reviewing of themes, defining and naming themes, and producing a report. Coding was first undertaken by the primary author (BCB) and was then shown to the senior author (DH) for critical appraisal and refinement. Following this, codes on similar topics were then collated, refined and grouped into a list of themes by both authors.

2.4 | Ethical considerations

This project was reviewed and approved by University College London Research Ethics Committee (Project ID Number: 14957/001).

3 | RESULTS

Table 1 outlines the themes and subthemes that were constructed in relation to the two aims of the study: (a) types of harm that exist from talking therapies with children and young people and (b) mechanisms for harm.

Two types of harm were identified by participants: 'Retraumatization' and 'Clinical Deterioration'. Additionally, four groups of mechanisms were identified as follows: 'Administrative factors', 'Relationship factors', 'Therapist factors' and 'Contextual factors'.

3.1 | Types of harm that can occur from talking therapies

All participants identified at least one type of harm that could occur from psychotherapy.

TABLE 1 Themes and subthemes in relation to clinicians' views of harm

Question	Themes	Subthemes
Types of harm	<i>Clinical deterioration</i> <i>Retraumatization</i>	
Mechanisms	Administrative factors Relationship factors Therapist factors Contextual factors	<i>Assessment processes</i> <i>Choosing the right intervention</i> <i>Child–Caregiver relationship</i> <i>Child–Therapist relationship</i> <i>Therapists' ability to reflect</i> <i>Insufficient support</i> <i>Lack of time</i> <i>Individualised focus of 1-1 therapy</i>

3.1.1 | Clinical deterioration

When asked about harm in psychotherapy, most participants highlighted clinical deterioration:

... [If] partway through therapy or by the end of the therapeutic process, a child is functioning less well, or feels less well than they were when they started.

(Participant 11)

Whilst getting worse in some circumstances was acknowledged as inevitable, it was felt that the worsening, or lack of progress with symptoms, could create a situation where children, young people or parents may not seek help in the future as therapy was seen as ineffective:

So many young people I've worked with, have had, 10 therapists? They've tried every modality under the sun, and will say 'quite frankly, it's bollocks', or, 'I never wanna go there again'.

(Participant 7)

3.1.2 | Retraumatization

As well as clinical deterioration, participants also mentioned retraumatization as a possible harmful effect of psychotherapy. This was prominent in clinicians who had background in trauma, abuse and neglect:

... You can run the risk of pushing something that actually the child or young person isn't ready to think about. And that can probably or could be potentially retraumatizing.

(Participant 8)

To mitigate against this, some participants outlined how support networks around the young person were critical. If support networks were not able to withstand or hold the treatment effects of trauma work, this could lead to the possibility of harm occurring outside of the clinical context:

Well, for trauma work, I would be looking at how containing the support network is around them. So, I wouldn't start working with a young person on a trauma if they didn't have a strong, stable, emotionally supportive environment that is going to be able to manage any emotional material that came up in between sessions.

(Participant 2)

3.2 | Mechanisms

3.2.1 | Theme I: Factors related to therapeutic administration

Many clinicians mentioned factors related to therapeutic administration which could result in harm. These focused on the assessment process, timing and choosing the right intervention.

Subtheme I: Assessment processes

Participants mentioned that a thorough assessment was important in helping mitigate against harm. During the first session(s) where assessment was undertaken, clinicians felt it was important to get to know the patients and their support networks, as this would create the set-up for subsequent meetings:

The first few sessions is finding out about, with their parents, finding out about their past and their story and feeling my way through how the family talks about it or doesn't talk about it, how the child responds to

ideas about their birthparents, or their birth sibling or the things that they've experienced.

(Participant 10)

Not undertaking a thorough and comprehensive assessment could result in harm as important information could be missed which the clinician needed to know about for formulation. For some, it was important to understand the clients' presenting difficulties from 'multiple perspectives' (Participant 5) and modalities so that they had a full and thorough understanding of the current situation.

Subtheme II: Choosing the right intervention

Towards the end of the assessment, clinicians discussed treatment options with the patient and family. Here, it was acknowledged that different types of interventions could treat a given presenting difficulty, but that client preferences and views should be taken into account. However, if the clinician was trained in a single therapeutic modality, the patient would end up receiving that intervention:

... single modality training of therapists, which means that if you go to see a play therapist, you're gonna get play therapy as your intervention [...] If you're gonna see a CBT therapist, you're gonna get CBT.

(Participant 5)

Ultimately, this may not be the right 'fit' and could result in the patient not getting better or dropping out.

Similar views were also posited more broadly in relation to types of treatment, with some clinicians querying whether talking therapies would be the best option when other types of support were available:

... There are lots of routes in to go to mental health and talking therapy is just often a tiny tiny part of it...I don't think we have a good idea of when it is [the best treatment option] and when it isn't.

(Participant 9)

Involving other professionals was seen as useful here, as well as discussing options with the patient.

3.2.2 | Theme II: Relationship factors

Around half of participants mentioned that harm could occur as a result of power dynamics, as the young person was in a weaker position of authority than both the clinician and their parent/guardian.

Subtheme I: Child-Caregiver relationship

Whilst clinicians highlighted that parents and guardians had an important role to play, some questioned whether harm could be caused as a result of children and young people being forced to attend therapy. As a result of this, they would not be assenting to treatment:

Children sometimes get pulled along and dragged along by the parents. So, I think that's more complicated, to get them to make a well-informed decision and to consent to things than it is for adults, so I think that increases the vulnerability.

(Participant 6)

This was highlighted by another clinician, who outlined the importance of making the young person feel comfortable to express themselves. Here, they acknowledged that the age difference between a young person and adult resulted in a power imbalance, which, if not acknowledged, meant a young person could be caused harm from being kept in treatment when they were no longer finding it helpful:

The ages create a very different power dynamic, where a child perhaps not knowing what to expect may stay within a therapy for longer with the belief that "the adults have made this decision this is something that I have to go along with", and they find it harder to speak up if they aren't finding the therapy helpful as well.

(Participant 7)

Subtheme II: Child-Therapist relationship

Like the above subtheme, power imbalances from the child and clinician dyad could also cause harm. Clinicians spoke about being in a position of power, where a child or young person may trust their judgement and expertise. If not managed correctly, this could result in unintentional coercion ending in harm:

I think they [children and young people] are potentially more vulnerable [...] because of that, there's more potential for coercion, for the therapist to kind of go "yeah come on let's do this" and the child being "okay". You know, they trust us, and they go along with it even if they're not ready.

(Participant 6)

For traumatic experiences, clinicians highlighted how this could be particularly harmful as these patients often entered therapy already disempowered and without a sense of control:

You can run the risk of pushing something that actually the child or young person isn't ready to think about. And that can probably or could be potentially retraumatising, but also can just be experienced as forceful [...] and I think that is particularly harmful for trauma victims who haven't had a say in what's happened to them.

(Participant 8)

In both instances described above, the need to develop a strong, secure and open therapeutic relationship was seen as important, and not doing so increased the potential for harm:

If they then feel that they can't even count on the adult who's there to help them, it kind of ... they lose hope I feel, so it's really important to try and create that hope, and to try and create that safe place... if you can't do that as a therapist I think you are causing harm just by the experience really.

(Participant 10)

3.2.3 | Theme III: Therapist factors

Subtheme: Therapists' ability to reflect

Participants mentioned that the inability of the therapist to reflect on if or why treatment was not working had the potential to result in harm:

I think one of the dangers is, well, is I know that some therapists and some psychologists don't tolerate easily or acknowledge when therapy isn't working.

(Participant 7).

Reflecting on when an approach was unsuccessful meant changes could be made, rather than harming the patient via continuing with ineffective treatment. Possible reasons for why reflecting may not happen included the therapist not having access to good supervision, as well as being burnt out due to being overstretched and time poor (see contextual factors). This was further highlighted by another clinician, who stated the importance of reflecting and being mindful of their thoughts, actions and behaviours when interacting with the patient:

I think that the most important thing, if you are really engaged with the person, and you're really working within the relationship, then I think it is important to be mindful at any stage throughout the therapeutic process, of things going wrong, the possibility of things being unhelpful, the possibility of my own perceptions and responses being counterproductive, mine as the therapist's, and putting that on the table.

(Participant 1)

Having open and honest conversations in this instance was thought to help mitigate this.

3.2.4 | Theme IV: Contextual factors

Three subthemes were identified as mechanisms for harm under contextual factors: insufficient support, lack of time, and the individualised focus of 1-1 therapy.

Subtheme I: Insufficient support

Many participants considered that when therapists do not have a supportive team or enough/good quality supervision, the risk of harm increases:

I think a lot of problems emerge for therapists when they don't have the adequate support structure within their organisations to do the work. So, if you have people who are working with families where, for example, young people are showing violence, they need very good clinical supervision that is adequate, that they can call on when needed, not just when provided, and by someone who is very familiar with the exact kind of work they're doing.

(Participant 1)

For other clinicians, insufficient organisational support, such as added pressure to see more patients, increased the risk of harm due to mistakes being made:

[There are] professionals in a context in an institution where they are so pushed, and so swamped to see people that they have burnt out and mistakes therefore happen.

(Participant 3)

Subtheme II: Lack of time

Clinicians thought there was a tension between wanting to allow the patient space to explore difficulties, whilst also knowing there were time limits imposed on how many times the patient could be seen:

There's little flexibility in the number of sessions you provide for instance, which is often the case of the NHS or even here, you can run the risk of pushing something that actually the child or young person isn't ready to think about.

(Participant 8)

As a result, this tension could result in the need to progress things faster than what would be optimal, or leaving the child open and exposed.

Participants also expressed that harm could be caused from discharging patients if they failed to attend a set number of sessions. They spoke of how, for some, it took time to build up trust to come to services. Discharging them too quickly for not attending treatment could play into the patients' feelings of abandonment and rejection:

And then also, some of the problems with, again due to pressures of CAMHS and resources, often if the young person doesn't attend a couple of sessions, they may be discharged, and there's so many people that have had a history of ongoing kind of rejection, from adults, or really difficult early life experiences, it

can sometimes take a long time to engage with these young people, so you might have a couple of missed appointments before you get them in.

(Participant 7)

This could lead to harm as their views became entrenched and meant they could be less likely to seek help in the future.

Subtheme III: Individualised focus of 1-1 therapy

A few clinicians outlined how the focus of one-on-one, individual therapy often drew on the notion that the problem was internally located in the young person. This could be damaging and harmful, creating the narrative of 'there's something wrong with me', 'I'm broken', 'I need fixing' (Participant 7). This was also echoed by another clinician, who felt that there was a general move within the clinical psychology profession to focus on the individual more than systemic approaches:

I think clinical psychology has very much entered into a medical model, and part of that is locating the disturbance almost entirely within, internal factors of the client, in service user, and losing awareness of external factors that we also need to consider, and have a more holistic understanding of what's going on.

(Participant 1)

4 | DISCUSSION

The aim of this study was to explore clinicians' perspectives on harm in psychotherapy when working with children and young people. Specifically, it set to explore what types of harm clinicians could identify, as well as mechanisms around how harm is caused. Each of these will be explored below.

4.1 | Types of harm

Clinicians identified two types of harm that could occur: clinical deterioration and retraumatisation. Clinical deterioration as a type of harm has previously been identified within the literature, both in quantitative studies and qualitative studies (Jonsson et al., 2016; Lilienfeld, 2007; Mohr, 1995; Schermuly-Haupt et al., 2018). Indeed, it has been suggested that this may be one of the most frequent side effects from talking therapies (Schermuly-Haupt et al., 2018). Deterioration in clinical samples involving children and young people has been shown to be around 7.7% for anxiety and 3.7% for depression (Edbrooke-Childs, Wolpert, Zamperoni, Napoleone, & Bear, 2018). Not all clinicians identified this as a type of harm which fits with the notion that additional factors need to be taken into account when making these judgements (Hardy et al., 2019; Linden, 2013). One reason for this is some patients do get worse

before they get better (Kramer & Stiles, 2015), and clinicians may view this as part of the therapeutic process.

Retraumatisation was the other type of harm identified. Unlike clinical deterioration, which could occur with any patient group, this is more nuanced and linked to the patient having to relive an identified trauma. None of the models reviewed in this paper address retraumatisation as an adverse event (Hardy et al., 2019; Lilienfeld, 2007; Linden, 2013). Thus, this may be considered a novel type of harm. Whilst there is overlap with similar concepts already identified, such as the emergence of new symptoms (Linden, 2013), this is different as it suggests a deliberate reliving of a pre-existing, buried emotion, which causes distress.

When compared to domains of the UE-ATR checklist, clinicians are not able to readily identify some of the categories identified. This appears to be particularly around events that occur outside of clinical settings. However, differences in terms of use, as well as not providing clinicians with a definition of harm at the beginning, such as in other research (Jonsson et al., 2016), may have contributed to these differences. Regardless, exploring clinicians' experience and notions of harm is relevant to enrich previous findings and identify different types of harm that could then be classified according to different criteria.

4.2 | Mechanisms of harm

In terms of how harm can be caused, four overarching categories were identified as follows: administrative factors, relationship factors, therapist factors and contextual factors.

Clinicians felt that in order to mitigate against harm, undertaking a thorough assessment was essential. This allowed the clinician to understand the presenting difficulties and formulate the best treatment plan. Additionally, selecting the 'right' treatment, which was congruent with the patients' values and preferences, was also considered important. Previous research around negative experiences has highlighted that clinicians should understand patients' psychosocial context to get the right fit (Hardy et al., 2019). Yet, children and young people frequently report being left out of conversations during appointments (Edbrooke-Childs, Calderon, Wolpert, & Fonagy, 2015; Hayes, Edbrooke-Childs, Town, Wolpert, & Midgley, 2019). This can result in an increased risk of drop out as the patient does not feel comfortable talking to the clinician, as well as not liking the structure and activity of the prescribed therapeutic modality (O'Keeffe, Martin, Target, & Midgley, 2019).

The therapist being in a dominant position over the patient has been previously highlighted as leading to harm. Findings here partially fit with previous research exploring clinicians' perceptions of negative events (Jonsson et al., 2016), as both highlight how a clinician, when overbearing, could lead to a compliant young person who continues treatment without expressing concerns or worries. Additionally, whilst the finding that parents too can dominate a conversation is not novel (Hayes, Edbrooke-Childs, Town, Wolpert, & Midgley, 2018), this is the first empirical study highlighting its role as

a mechanism of harm. To counteract power dynamics, clinicians may wish to engage in shared decision-making and make sure they have time alone with the young person (Hayes, Fleming, & Wolpert, 2015).

The reflective capacity of the clinician was also deemed to be an important factor. One aspect of this focused on the clinician reflecting on their relationship with the patient, with harm caused when the therapist was not being mindful. The therapeutic relationship is an important aspect in determining outcomes (Shirk, Karver, & Brown, 2011). In terms of harm, clinicians have highlighted this poor therapeutic alliance as a contributing risk factor (Jonsson et al., 2016). The other aspect around reflective capacity focused on the clinician not recognising when the treatment was ineffective, which could cause harm if the treatment was continued. The ability of clinicians to predict treatment outcome and deterioration is poor and to overcome this, it has been suggested that regular monitoring and real-time change are needed (Lambert, 2011).

Wider contextual factors focused on insufficient support, a lack of time, and the individualised focus of therapy. Insufficient support focused on not getting good quality supervision, as well as clinicians having high workloads. Inadequate supervision and its effect on harm have previously been documented and are potentially more harmful when the therapist is inexperienced or has a complex case (Hardy et al., 2019). However, even when supervision is available, it may not be used appropriately for issues around harm due to time constraints (Hardy et al., 2019).

Linked with insufficient support was also a lack of time. Here, clinicians highlighted the tension between not wanting to rush the patient into opening up, versus knowing there was only a set number of sessions they could provide. They further highlighted that when patients did not attend appointments, there was a pressure to discharge them so they could see another. Discharging too early could be harmful as these patients may take time to build up trust with a clinician. Letting them go would confirm the patients' experiences of being 'broken' or 'being impossible to help', leading them to not seeking help in the future. Factors around time have previously been outlined as contributing to harm, with an emphasis on building a strong trusting therapeutic relationship, and the need to balance time against service demands (Hardy et al., 2019). With time being a precious commodity and demand unlikely to fall, open and honest conversations at the start of therapy about length and what can be achieved may be a necessary compromise (Hayes et al., 2015), as well as an expedited route back for individuals who have previously failed to attend.

The last contextual factor focused on how certain types of therapeutic approaches could lead to too much focus on the patient, creating a narrative that there was something wrong with them. Stigmatisation is one of the unwanted events outlined in the Linden (2013) model, and within empirical papers, patients cite feelings of failure as an outcome related to ineffective therapy (Hardy et al., 2019). To counteract this, clinicians may wish to explore the biopsychosocial approach with patients during assessment, as well as the precipitating, perpetuating, predisposing and present factors around their mental health difficulty. Therapeutic modality and

patient values are also an important consideration, as congruence between these may help mitigate negative patient experiences.

4.3 | Limitations of the study

Whilst this study contributes to an under-developed field of research, limitations exist. In particular, there was a skew towards clinicians who worked in trauma, as well as an integrative framework, meaning some types of mechanisms may have been missed, or over-represented in these findings. Another further limitation is the sample size, which might have meant not reaching saturation of themes.

A further aspect to consider is that in terms of the types of harm involved or the mechanisms by which it may happen, some types of harm may fall outside the awareness of clinicians, due to relational or other factors such as not adhering to therapeutic principles. Indeed, some clinicians spoke about theoretical types of harm that could occur, rather than harm that had occurred in their practice or harm they knew patients who had come to them had previously experienced.

4.4 | Further lines of research

Further lines of research may include exploring harm with clinicians working in specific settings. This is relevant given the finding of 're-traumatisation' as a type of harm highlighted by those working in that field. Additionally, exploration with children, young people and parents around harm is also needed, particularly as there is a lack of congruence between them around the types of mental health difficulties and treatment options (Yeh & Weisz, 2001). This may allow for the identification of harmful experiences which remain out of the awareness of particular stakeholder groups, especially since patients' and therapists' understanding of negative effects are divergent (Curran et al., 2019).

4.5 | Implications

- Clinicians should be aware of the possibility of different types of harm happening and how this could impact on their own practice. This is especially true given that clinicians sometimes struggle to identify when things go wrong (Lambert, 2011).
- The importance of assessment has been highlighted, as well as the need for clinicians to have good quality, regular supervision.
- Including patients in care and treatment decisions can help mitigate against harm, as the option will be aligned with the values and help create an open and honest relationship.
- Health institutions should consider utilising data to understand when patients are deteriorating and the possible reasons for this.

ORCID

Dan Hayes  <https://orcid.org/0000-0003-4948-3333>

REFERENCES

- Barlow, D. H. (2010). Negative effects from psychological treatments: A perspective. *American Psychologist*, 65(1), 13–20. <https://doi.org/10.1037/a0015643>
- Blease, C. R., Lilienfeld, S. O., & Kelley, J. M. (2016). Evidence-based practice and psychological treatments: The imperatives of informed consent. *Frontiers in Psychology*, 7, 1–5. <https://doi.org/10.3389/fpsyg.2016.01170>
- Bowie, C., McLeod, J., & McLeod, J. (2016). 'It was almost like the opposite of what I needed': A qualitative exploration of client experiences of unhelpful therapy. *Counselling and Psychotherapy Research*, 16(2), 79–87. <https://doi.org/10.1002/capr.12066>
- Braun, V., & Clarke, V. (2006). Using thematic analysis in psychology. *Qualitative Research in Psychology*, 3(2), 77–101. <https://doi.org/10.1191/1478088706qp063oa>
- Bystedt, S., Rozental, A., Andersson, G., Boettcher, J., & Carlbring, P. (2014). Clinicians' perspectives on negative effects of psychological treatments. *Cognitive Behaviour Therapy*, 43(4), 319–331. <https://doi.org/10.1080/16506073.2014.939593>
- Curran, J., Parry, G. D., Hardy, G. E., Darling, J., Mason, A. M., & Chambers, E. (2019). How does therapy harm? A model of adverse process using task analysis in the meta-synthesis of service users' experience. *Frontiers in Psychology*, 10, 347. <https://doi.org/10.3389/fpsyg.2019.00347>
- Dimidjian, S., & Hollon, S. D. (2010). How would we know if psychotherapy were harmful? *American Psychologist*, 65(1), 21–33. <https://doi.org/10.1037/a0017299>
- Duggan, C., Parry, G., McMurran, M., Davidson, K., & Dennis, J. (2014). The recording of adverse events from psychological treatments in clinical trials: Evidence from a review of NIHR-funded trials. *Trials*, 15(1), 335. <https://doi.org/10.1186/1745-6215-15-335>
- Edbrooke-Childs, J., Calderon, A., Wolpert, M., & Fonagy, P. (2015). *Children and young people's improving access to psychological therapies rapid internal audit: National report*. London.
- Edbrooke-Childs, J., Wolpert, M., Zamperoni, V., Napoleone, E., & Bear, H. (2018). Evaluation of reliable improvement rates in depression and anxiety at the end of treatment in adolescents. *BJPsych Open*, 4(4), 250–255. <https://doi.org/10.1192/bjo.2018.31>
- Hardy, G. E., Bishop-Edwards, L., Chambers, E., Connell, J., Dent-Brown, K., Kothari, G., ... Parry, G. D. (2019). Risk factors for negative experiences during psychotherapy. *Psychotherapy Research*, 29(3), 403–414. <https://doi.org/10.1080/10503307.2017.1393575>
- Hayes, D., Edbrooke-Childs, J., Town, R., Wolpert, M., & Midgley, N. (2018). Barriers and facilitators to shared decision making in child and youth mental health: Clinician perspectives using the Theoretical Domains Framework. *European Child and Adolescent Psychiatry*, 28(5), 655–666
- Hayes, D., Edbrooke-Childs, J., Town, R., Wolpert, M., & Midgley, N. (2019). Barriers and facilitators to shared decision-making in child and youth mental health: Exploring young person and parent perspectives using the Theoretical Domains Framework. *Counselling and Psychotherapy Research*, 20(1), 57–67. <https://doi.org/10.1002/capr.12257>
- Hayes, D., Fleming, I., & Wolpert, M. (2015). Developing safe care in mental health for children and young people: Drawing on UK experience for solutions to an under-recognised problem. *Current Treatment Options in Pediatrics*, 1(4), 309–319. <https://doi.org/10.1007/s40746-015-0037-1>
- Jonsson, U., Johanson, J., Nilsson, E., & Lindblad, F. (2016). Adverse effects of psychological therapy: An exploratory study of practitioners' experiences from child and adolescent psychiatry. *Clinical Child Psychology and Psychiatry*, 21(3), 432–446. <https://doi.org/10.1177/1359104515614072>
- Klatte, R., Strauss, B., Flückiger, C., & Rosendahl, J. (2018). Adverse effects of psychotherapy: Protocol for a systematic review and meta-analysis. *Systematic Reviews*, 7(1), 1–8. <https://doi.org/10.1186/s13643-018-0802-x>
- Kramer, U., & Stiles, W. B. (2015). The responsiveness problem in psychotherapy: A review of proposed solutions. *Clinical Psychology, Science and Practice*, 22, 277–295. <https://doi.org/10.1111/cpsp.12107>
- Lambert, M. J. (2011). What have we learned about treatment failure in empirically supported treatments? Some suggestions for practice. *Cognitive and Behavioral Practice*, 18(3), 413–420. <https://doi.org/10.1016/j.cbpra.2011.02.002>
- Lilienfeld, S. O. (2007). Psychological treatments that cause harm. *Perspectives on Psychological Science*, 2(1), 53–70. <https://doi.org/10.1111/j.1745-6916.2007.00029.x>
- Linden, M. (2013). How to define, find and classify side effects in psychotherapy: From unwanted events to adverse treatment reactions. *Clinical Psychology and Psychotherapy*, 20(4), 286–296. <https://doi.org/10.1002/cpp.1765>
- Mercer, J. (2017). Evidence of potentially harmful psychological treatments for children and adolescents. *Child and Adolescent Social Work Journal*, 34(2), 107–125. <https://doi.org/10.1007/s10560-016-0480-2>
- Mohr, D. C. (1995). Negative outcome in psychotherapy: A critical review. *Clinical Psychology: Science and Practice*, 2, 1–27. <https://doi.org/10.1111/j.1468-2850.1995.tb00022.x>
- Moritz, S., Fieker, M., Hottenrott, B., Seeralan, T., Cludius, B., Kolbeck, K., ... Nestoriuc, Y. (2015). No pain, no gain? Adverse effects of psychotherapy in obsessive-compulsive disorder and its relationship to treatment gains. *Journal of Obsessive-Compulsive and Related Disorders*, 5, 61–66. <https://doi.org/10.1016/j.jocrd.2015.02.002>
- Moritz, S., Nestoriuc, Y., Rief, W., Klein, J. P., Jelinek, L., & Peth, J. (2018). It can't hurt, right? Adverse effects of psychotherapy in patients with depression. *European Archives of Psychiatry and Clinical Neuroscience*, 269(5), 577–586. <https://doi.org/10.1007/s00406-018-0931-1>
- O'Keeffe, S., Martin, P., Target, M., & Midgley, N. (2019). 'I just stopped going': A mixed methods investigation into types of therapy dropout in adolescents with depression. *Frontiers in Psychology*, 10, 75. <https://doi.org/10.3389/fpsyg.2019.00075>
- Parry, G. D., Crawford, M. J., & Duggan, C. (2016). Iatrogenic harm from psychological therapies-time to move on. *British Journal of Psychiatry*, 208(3), 210–212. <https://doi.org/10.1192/bjp.bp.115.163618>
- QSR International (2018). NVivo qualitative data analysis software. QSR International Pty Ltd.
- Schermuly-Haupt, M. L., Linden, M., & Rush, A. J. (2018). Unwanted events and side effects in cognitive behavior therapy. *Cognitive Therapy and Research*, 42(3), 219–229. <https://doi.org/10.1007/s10608-018-9904-y>
- Shirk, S. R., Karver, M. S., & Brown, R. (2011). The alliance in child and adolescent psychotherapy. *Psychotherapy*, 48(1), 17. <https://doi.org/10.1037/a0022181>
- Vaughan, B., Goldstein, M. H., Alikakos, M., Cohen, L. J., & Serby, M. J. (2014). Frequency of reporting of adverse events in randomized controlled trials of psychotherapy vs. psychopharmacotherapy. *Comprehensive Psychiatry*, 55(4), 849–855. <https://doi.org/10.1016/j.comppsy.2014.01.001>
- Yeh, M., & Weisz, J. R. (2001). Why are we here at the clinic? Parent-child (dis)agreement on referral problems at outpatient treatment entry. *Journal of Consulting and Clinical Psychology*, 69(6), 1018–1025. <https://doi.org/10.1037/0022-006X.69.6.1018>

AUTHOR BIOGRAPHIES

Ms Bárbara Castro Batic did her undergraduate studies in Psychology, at Pontificia Universidad Católica de Chile. She is currently working as a psychotherapist in Chile, and has a MSc in Psychoanalytic Developmental Psychology. She is interested in the public mental health field and in psychoanalytic theory.

Dr Daniel Hayes is a Senior Research Supervisor on the MSc Psychoanalytic Developmental Psychology at University College London, as well as Research Fellow at the Evidence Based Practice Unit. His research interests include: shared decision making with children and young people, developing and evaluating community interventions, and safe care in mental health.

How to cite this article: Castro Batic B, Hayes D. Exploring harm in psychotherapy: Perspectives of clinicians working with children and young people. *Couns Psychother Res*. 2020;20:647–656. <https://doi.org/10.1002/capr.12347>

APPENDIX A

Exploring clinicians' perspectives on harm in psychotherapy with children and adolescents

- When I mention the word "harm", in the context of psychotherapy, what comes to mind?
- In your opinion, what type of risks are involved in psychotherapy, if any?
- Are there any risks for the clinician as well?
- Do you think children or adolescents can be more vulnerable to harm in psychotherapy than adults? If so, why? If not, why?
- What aspects are necessary to consider when making a decision about something being potentially harmful?