

The Urinal Problem

Evangelos Kranakis¹ and Danny Krizanc²

¹ School of Computer Science, Carleton University
K1S 5B6, Ottawa, Ontario, Canada

² Department of Mathematics and Computer Science
Wesleyan University, Middletown CT 06459, USA

Abstract. A man walks into a men’s room and observes n empty urinals. Which urinal should he pick so as to maximize his chances of maintaining privacy, i.e., minimize the chance that someone will occupy a urinal beside him? In this paper, we attempt to answer this question under a variety of models for standard men’s room behavior. Our results suggest that for the most part one should probably choose the urinal furthest from the door (with some interesting exceptions). We also suggest a number of variations on the problem that lead to many open problems.

1 Introduction

The question of digital privacy and how to protect it has a long history of study in computer science and it raises a number of interesting algorithmic (as well as other) research problems[7]. But it is also the case that algorithmic issues may arise when considering physical privacy. A particular instance of this occurs when one considers the use of a public men’s room. The standard design of a men’s room contains a number of urinals which are generally located along one wall with neighboring stations in full view of each other (Figure 1). (Although recently there has been a trend to place dividers between adjoining positions [1]. For a history of privacy concerns in restroom design see [15].) In order to obtain some amount of privacy while vacating one’s bladder it is desirable to have a urinal such that its neighboring positions are unoccupied¹.

This leads to the following algorithmic question: A man walks into a men’s room and observes n empty urinals. Which urinal should he pick so as to maximize his chances of maintaining privacy, i.e., minimize the chance that someone will occupy a urinal beside him? One’s intuition might suggest that choosing one of the end urinals is the best choice. It turns out the answer depends on many

¹ We hope that by focusing on the men’s room version of the problem we are not leaving out our female readers. We chose this version since (1) we understand (not from personal experience) that most women’s rooms contain only water closets which already provide some amount of privacy beyond that afforded by urinals and (2) we are more familiar with the behavior of men in a public restroom situation which we shall see is an important aspect of our study. It has been pointed out to us that women may also use urinals [2] and should this become a more prevalent behavior our results may directly interest our female readers.



Fig. 1. Standard men’s room urinal arrangement

things but in particular on how one models the behavior of the men who enter later. In this paper, we consider a variety of models for men’s room behavior and attempt to develop strategies for maintaining privacy under each. The results of this preliminary study suggest that for the most part one’s intuition is correct (the ends are best) with some interesting exceptions. We also examine a number of variations on the problem that might lead to a better understanding of physical privacy concerns and which suggest many interesting open questions.

2 Related Work

As far as we know the problem we consider has not been studied before. If one is interested in finding the most private position to place an object in the sense of it being furthest from all other occupied positions in some space this can be formulated as a version of the well-studied *Obnoxious Facility Location Problem*. (See [13] for a survey of results in this area.) But in our case this question is trivially solved by taking the position at the mid-point of the largest gap of urinals (i.e., contiguous sequence of empty urinals) and it does not really answer our question. The question of privacy in querying sensors in a sensor network also shares some aspects of our question but again the concerns turn out to be different [16].

Most closely related to our question (at least for the case where the men behave randomly) seems to be the following “unfriendly seating arrangement” problem posed by Freedman and Shepp [4]: There are n seats in a row at a

luncheonette and people sit down one at a time at random. They are unfriendly and so never sit next to one another (no moving over). What is the expected number of persons to sit down? Solutions to this problem were provided by Friedman, Rothman and MacKenzie [5,12] who show that as n tends to infinity the expected fraction of the seats that are occupied goes to $\frac{1}{2} - \frac{1}{2e^2}$. (For a nice exposition of this and related problems see [3].) Georgihou et al. [6] consider the following generalization of this problem (the unfriendly theatre seating arrangement problem): People arrive one at a time to a theatre consisting of m rows of length n . Being unfriendly they choose seats at random so that no one is in front of them, behind them or to either side. What is the expected number of people in the theatre when it becomes full, i.e., it can not accommodate any more unfriendly people? They give bounds on the fraction of the theatre that is full (in the limit) for all m and show that for $m = 2$ this limit is $\frac{1}{2} - \frac{1}{4e}$.

While perhaps related to our problem in name only, our study was at least partially inspired by Knuth’s Toilet Paper problem [11]. Don Knuth relates that the toilet paper dispensers in the Stanford Computer Science public restrooms are designed to hold two rolls of tissues either of which is available for use. This led him to consider the following problem: A toilet stall contains two rolls of toilet paper of n sheets each. The stall is used by people of two types: *big choosers* and *little choosers*. They arrive to use the toilet randomly and independently, the former with probability p and the latter with probability $1 - p$. Big (respectively, little) choosers select exactly one sheet of paper from the roll with the most (respectively, least) number of sheets. What is the expected number of toilet sheets remaining just after one of the two rolls has emptied, defined to be the residue $R_n(p)$?

Knuth used combinatorial techniques to prove that for fixed p and r , which satisfy the condition $4p(1 - p) < r < 1$, we have that

$$E[R_n(p)] = \begin{cases} \frac{p}{2^{p-1}} + O(r^n) & \text{if } p > 1/2 \\ 2\sqrt{\frac{p}{\pi}} - \frac{1}{4}\frac{1}{\sqrt{n\pi}} + O(n^{-3/2}) & \text{if } p = 1/2 \\ \frac{1-2p}{1-p}n + \frac{p}{1-2p} + O(r^n) & \text{if } p < 1/2 \end{cases}$$

as $n \rightarrow \infty$ where the constants implied by the O notation depend on p, r but not on n . A high probability version of this result was recently used to prove bounds on the time required for routing on a Markovian grid [9].

3 The Urinal Problem

For our initial study we assume that you are the first man to enter and all of the urinals are available. We will discuss the case where some urinals are occupied below. Further we assume that men enter one at a time and depending on one of the strategies outlined below they make a choice of a urinal that provides privacy (i.e., is unoccupied on either side) if possible. If this is not possible

then we assume the *random* strategy for choosing a urinal where one chooses uniformly at random among all unoccupied urinals. We note that other models of how the remaining urinals are filled are possible and make some remarks on some of them below. Finally, we assume that men enter at a constant rate of one man every time unit and stay at their position until you have finished.

For each of the scenarios we consider, we are interested in maximizing the expected time until your privacy will be violated, i.e., a man occupies one of the urinals beside yours. We call a set of the occupied urinals a *configuration*. We call a configuration *saturated* if the next man entering is forced to violate at least one person's privacy. It is fairly clear that, under the assumptions above, the time until your privacy is violated can be divided into two phases: the time until saturation is reached and time until a man violates your privacy once the configuration is saturated. As such we divide our analysis for each of the cases below into two parts: the time until saturation and the time until privacy violation once saturation has been achieved.

3.1 Lazy Behavior

In this model one chooses the lowest number unoccupied urinal that provides privacy, i.e., we assume that a urinal's distance from the door of the men's room is directly proportional to its number and that men, being naturally lazy, will always choose the first empty private spot. The analysis of the first phase divides into two cases depending upon whether n is even or odd.

If n is even and you choose an odd numbered position then clearly the saturated configuration will consist of all of the odd positions and will contain $n/2$ occupied urinals. If you choose an even numbered position, $2k$, then the saturated configuration consists of the $k - 1$ odd positions before you (excluding $2k - 1$) and the $n/2 - k$ even positions after you. Again there are $n/2$ occupied urinals at saturation. In the odd case, you are better off choosing an odd position as in that case the saturated configuration consists of all of the odd positions and contains $(n + 1)/2$ men versus $(n - 1)/2$ men in case you choose an even position.

In either case, choosing any odd position yields $\lceil n/2 \rceil$ men in the saturated configuration. At this point, if the remaining positions are filled randomly, there is a distinct advantage to picking one of the positions at the end. The number of men that enter before picking a urinal beside you follows a negative hypergeometric distribution with parameters $N = n - \lceil n/2 \rceil$, a where a is the number of positions available beside you, i.e., $a = 1$ if you occupy position 1 or n and $a = 2$ otherwise. The expectation of such a random variable is given by $(N + 1)/(a + 1)$ so we get the expected time until your privacy is violated is

$$\lceil n/2 \rceil + \frac{\lfloor n/2 \rfloor}{2}$$

if you choose positions 1 or n and

$$\lceil n/2 \rceil + \frac{\lfloor n/2 \rfloor}{3}$$

otherwise. Clearly, choosing an end position is to your advantage for all n . (We discuss the case where urinals are filled in a lazy fashion after saturation is reached below.)

3.2 Cooperative Behavior

In this model, all men cooperate to ensure that the configuration becomes saturated at the last possible moment, by each choosing a position that guarantees the maximum number of men have full privacy for as long as possible. It is easy to establish that this case is very similar to the above in that as long as you choose an odd position the saturated configuration will contain $\lceil n/2 \rceil$ men and the best overall choice turns out to be either end.

3.3 Maximize Your Distance Behavior

In this model we assume that a new entrant to the washroom will choose the urinal which maximizes his distance to any of the current occupants of urinals. If more than one such urinal exists then a random one among them is chosen. If no urinal with privacy exists then a random urinal is chosen.

First we observe that by symmetry the number and positions of the men when saturation is first reached are independent of the random choices made among equidistant positions and only depends on the choice that you make as the first man. Let $A(n, i)$ be number of men in the saturated state if the first man chooses position $i, i = 1, \dots, n$. Let $B(n)$ be the number of men in such a saturated state assuming positions 1 and n are filled (not including the men at urinals 1 and n).

Assume $n > 3$. (The cases $1 \leq n \leq 3$ are straightforward to analyze.) Observe that if the first man chooses position 1 (respectively, n) then the second one will choose n (respectively, 1) and the saturated configuration will contain $2 + B(n)$ men. If the first man chooses 2 (respectively, $n - 1$) the second will choose n (respectively, 1). If the first man chooses some position i between 3 and $n - 2$ then the second man will choose the further of 1 and n and eventually someone will choose the other. This yields the following equation for $A(n, i)$:

$$A(n, i) = \begin{cases} 2 + B(n) & i = 1, n \\ 2 + B(n - 1) & i = 2, n - 1 \\ 3 + B(i) + B(n - i + 1) & 3 \leq i \leq n - 2. \end{cases}$$

Further observe that in any gap of $k > 2$ unoccupied urinals with the urinals at either end occupied, the first of the unoccupied positions to be occupied will be the middle one (or one of the two middle ones if k is even). This yields:

$$B(n) = \begin{cases} 0 & n \leq 4 \\ 1 + B(\lfloor \frac{i+1}{2} \rfloor) + B(\lceil \frac{i+1}{2} \rceil) & n > 4. \end{cases}$$

It is relatively straightforward to establish the following closed form solution for $B(n)$:

$$B(n) = 2^{\lfloor \log(n-1) \rfloor - 1} + (\lfloor \frac{n-1}{2^{\lfloor \log(n-1) \rfloor - 1}} \rfloor - 2)(n - 3 \cdot 2^{\lfloor \log(n-1) \rfloor - 1} - 1) - 1$$

which in turn allows us to compute $A(n, i)$ for any n, i . Unfortunately, we were not able to establish a closed form expression for $\max_i A(n, i)$ or show that it can be computed any faster than evaluating $A(n, i)$ for all i . Note that this is a pseudo-polynomial time algorithm as the input is just n . (See the discussion below concerning different initial configurations for more on the analysis of this algorithm.) While the function is reasonably well-behaved it does vary widely for some values of n between the extremes of $\lceil n/3 \rceil$ and $\lceil n/2 \rceil$.

If one was just concerned about maximizing the time until saturation occurs it turns out that end positions are not optimal. In fact, for some values of n position 1 is the worst choice in this regard. For example, for n of the form $3 \cdot 2^k + 1$ for some $k > 0$, $A(n, 1) = \lceil n/3 \rceil$ while $A(n, \lceil n/3 \rceil) = \lceil n/2 \rceil$.

Fortunately, one can show that once again the end positions have an advantage when we assume that the remaining positions are filled in random order. It is fairly easy to show that $A(n, i) \geq \lceil n/3 \rceil$ for all n (at least one of every three urinals must be occupied) and that $A(n, i) \leq \lceil n/2 \rceil$ (at most one of every pair may be occupied) for all n and i and since

$$\lceil n/3 \rceil + \frac{n - \lceil n/3 \rceil}{2} > \lceil n/2 \rceil + \frac{\lfloor n/2 \rfloor}{3}$$

we see that urinal number 1 (and n) is always (by a very slight margin in some cases, e.g., for n of the form $3 \cdot 2^k + 1$ for some $k > 0$ the difference is $1/6$) the optimal choice.

3.4 Random Behavior

In this model we assume that when a man enters if the configuration is not saturated he chooses uniformly at random among all positions that provide him with privacy on both sides and that once saturation is reached he chooses uniformly among all available positions. If the first man chooses randomly like everyone else then the question of how many men there are at the time of saturation corresponds to the unfriendly seating arrangement problem discussed above. In our case we are interested in computing whether it makes a difference which position you choose as the first man. And indeed it does. Consider the case $n = 5$. If position 1 is chosen it is easy to calculate that the expected number of men at saturation is 2.67 but if you choose position 3 you are guaranteed 3 men at saturation so it would appear that position 3 is better. But as above it is during the filling phase that the advantage of an end takes over. In this example, position 1 expects 4.33 men (including himself) before his privacy is violated versus the situation for position 3 where the fourth man will certainly violate his privacy.

Let $E(n, i)$ be the expected number of men entering before your privacy is violated assuming you choose position i . Let $p_{i,k}$ be the probability that if you choose position i the resulting saturated configuration has k men in it. Let $q_{j,k,\ell}$ be the probability that, if there are k spots available after saturation and $\ell = 1, 2$

depending on whether you are at an end or not (indicating the number of positions you have available beside you), your privacy is violated by the j th man ($j = 1, \dots, n - k$) entering after saturation. As above we note that a saturated configuration has between $\lceil n/3 \rceil$ and $\lceil n/2 \rceil$ positions taken and so we can calculate:

$$\begin{aligned}
 E(n, i) &= \sum_{s=\lceil n/3 \rceil+1}^n s \sum_{j+k=s, j>0, \lceil n/3 \rceil \leq k \leq \lceil n/2 \rceil} q_{j,n-k,\ell} p_{i,k} \\
 &= \sum_{k=\lceil n/3 \rceil}^{\lceil n/2 \rceil} (k + \sum_{j=1}^{n-k} j \cdot q_{j,n-k,\ell}) p_{i,k}.
 \end{aligned}$$

Using the fact that that $q_{j,k,\ell}$ is negative hypergeometrically distributed we get:

$$E(n, i) = \begin{cases} \frac{n+1}{2} + (1/2) \sum_{k=\lceil n/3 \rceil}^{\lceil n/2 \rceil} k p_{i,k} & i = 1, n \\ \frac{n+1}{3} + (2/3) \sum_{k=\lceil n/3 \rceil}^{\lceil n/2 \rceil} k p_{i,k} & i = 2, \dots, n - 1. \end{cases}$$

Let $F(n, i) = \sum_{k=\lceil n/3 \rceil}^{\lceil n/2 \rceil} k p_{i,k}$. Note that $F(n, i)$ is the expected number of urinals occupied at saturation if position i is chosen first. But it is easy to see that

$$F(n, i) = 1 + \begin{cases} F(n - 2) & i = 1, n \\ F(n - 3) & i = 2, n - 1 \\ F(i - 2) + F(n - i - 1) & n = 3, \dots, n - 2 \end{cases}$$

where $F(n) = \sum_{i=0}^{n-1} \frac{(-2)^i (n-1)}{(i+1)!}$ is the expected number of urinals at saturation if all choices are random established in [5,12].

While $F(n)$ is monotonic as we saw above for the case $n = 5$ it is not the case that $F(n, 1) > F(n, i)$ for all i . On the other hand, it is fairly straightforward to show that $F(n, 1) + 1 > F(n, i)$ and $F(n, i) \leq \lceil n/2 \rceil$ for all i and n and from this conclude (from the formula for $E(n, i)$ above) that once again positions 1 (and n) are the optimal choice.

4 Variations on Our Theme

4.1 Different Filling Strategies

The above analysis assumes once saturation is reached the remaining positions are filled uniformly at random. This may not be realistic as one could imagine other approaches.

Consider, for example, a *lazy* filling strategy whereby a new entrant, finding the configuration saturated, chooses the first available urinal. We say a position is *fully-private* if the urinals on either side of it (if they exist) are unoccupied. A non-end position is *semi-private* if only one of its neighbors is occupied. A position is *non-private* if it is not fully private or semi-private. We say a man's

behavior is *fully-private first* if a man always chooses a fully-private position if available. All of the behaviors we have been considering are fully-private first.

It is fairly straightforward to show that for any behavior that follows a fully-private first initial strategy and a lazy filling strategy, the optimal choice is always position n as in this case all urinals except $n - 1$ must be used before your privacy is violated.

An interesting situation arises in the case where semi-private positions are preferred to non-private positions when filling. The expected time until your privacy is violated now depends on how many semi-private positions versus non-private positions exist at saturation and what types of gaps exist beside you (one or two unoccupied positions). Given this information the expected time of your privacy can be calculated using the appropriate negative hypergeometric distribution.

Consider what happens in the case of the distance maximizing behavior above if (1) in the initial phase when choosing between two equi-distant alternatives one always chooses the lower numbered urinal (i.e., the lazy choice) and (2) after reaching saturation the men choose randomly first among semi-private positions and only after they are filled they choose randomly among non-private positions. In this case, the advantage held by the end positions disappears! If $n = 8$ the expected time until privacy is violated for position 1 is 5 versus 6.33 if you choose position 3. If $n = 21$ the expected length of your privacy for position 21 is 14.5 versus 15 if you choose position 3. In fact, one can construct infinitely many n for which position 1 (or position n) is not the best choice. ($n = 8$ is the smallest case for position 1 and $n = 6$ the smallest for the last position.) But in each of these cases, we find that if the first urinal is not optimal, the last is, and vice versa. We conjecture that under the behavior above it will always be the case that either position 1 or position n will be optimal. Further we conjecture that if instead of making a lazy choice for equidistant positions in the first phase but make a random choice, the advantage for both position 1 and position n remains. If your filling strategy is to always fill a randomly chosen gap of size 2 in order to get a semi-private position (if one exists) in a lazy manner, then one can show that position 1 is no longer optimal (already it fails for $n = 4$) but position n is always optimal. For the random behavior and for any of the filling strategies considered above one can show that position n remains optimal.

4.2 Non-empty Initial Configuration

More often than not when you enter a public men's room in a busy area the configuration you observe is not the empty one. What should one do in this case?

A rule of thumb that follows from the discussion above is it likely the case that if one of the end positions is open and its single neighbor is unoccupied, you should choose that position. Note that this is not always the case! If we are in a situation where one of the ends is available but choosing it will force a gap of size 2 to form beside it, it may be the case that, if semi-private positions are preferred after saturation, choosing the end is not a good idea.

In general, the only way we know of to be certain you make the best choice is to compute for each position the expected length of privacy using the formulas provided above applying them to the subproblems created by the configuration. For example, if one is analyzing the case where maximizing the distance behavior is assumed, then one can compute the time to saturation by adding the times resulting from each of the gaps in the configuration using the formula $B(n)$ (or an appropriate modification in the case where one end of a gap is position 1 or n). After saturation, the filling time can be computed using the appropriate negative hypergeometric distribution. If the configuration is presented as $\{0, 1\}$ vector then this results in a $O(n^2)$ time algorithm for finding the best urinal to choose. If the configuration is given as a list of positions then in general the algorithm is only pseudo-polynomial. We conjecture there is a polynomial time algorithm for each of the behaviors and filling strategies discussed above but have not been able to determine one for all cases.

4.3 Dynamic Situation

Perhaps the least realistic assumptions we have made above are (1) the arrival rate of men is constant and (2) the men stay indefinitely once they arrive rather than leaving once they have completed their business. It is fairly evident that both the arrivals and service times of the men are complex random processes. As a first attempt, the use of queueing theory to model this aspect of the problem seems warranted.

The simplest assumption in this regard would be that the arrival times form a Poisson process and the time required to relieve oneself is exponential distributed. As there are n urinals, this leads to what is termed an M/M/ n queue in the standard queueing theory terminology [10,14]. Given $1/\lambda$, the mean time between arrivals, and $1/\mu$, the mean service time, it is straightforward to calculate such quantities as the expected number of “customers” in the system (either being serviced or waiting for a urinal to open up in case all are full), the probability that k urinals are free, etc. and use this information to help devise a strategy for picking a urinal.

For example, if $\frac{\lambda}{\mu} < n/2$ then you are in a region where one expects that saturation will not be reached during your visit (or even if it is the number of filled positions after saturation will be small.) In this case, if one assumes the maximizing distance behavior above then the advantage of the ends disappears. One should use the algorithm above to estimate the position that prolongs the time until saturation the longest. Generally, this will be somewhere in the largest gap of urinals and not necessarily at the end. (Again, if we assume a configuration is represented by a $\{0, 1\}$ vector this will be a polynomial time algorithm.) On the other hand, if one assumes random behavior then for most configurations (ones with a sufficiently large gap) a search using a Python program suggests the best choice is one that leaves three empty urinals between you and either the beginning or the end of the largest gap. This appears to insure that the time to saturation in that gap will be maximized.

The above analysis assumes you have an good estimate of λ and μ which seems unlikely². Without this information one can make a maximum likelihood estimate of $\frac{\lambda}{\mu}$ using the number of men in the configuration upon your arrival. If the current configuration is not saturated you may as well assume that this is the steady-state and follow the guidelines described in the last paragraph for the case $\frac{\lambda}{\mu} < n/2$. On the other hand if the configuration is saturated then you have no choice but to pick a semi-private or even non-private position. It might be interesting to investigate whether one such spot is better than another for increasing your privacy. This would seem to depend on having an estimate of the time remaining to service individuals currently in the configuration but perhaps not.

4.4 Game-Theoretic Formulation

Another objection to the approach taken above is that those arriving after the first arrival are limited to following a particular behavior. In reality they are also interested in maximizing their private time. How might considerations of more general strategies effect your choice? If the second man to enter is going to maximize his privacy based upon your choice how does that effect your choice as first. If k men enter simultaneously, is there a mechanism by which they might be led to choose a configuration that maximizes the average or maxmin privacy?

These questions suggest that our problem is in reality some sort of game played by multiple players in rounds and the tools of game theory might be applied to establish better strategies. We leave it up to experts in algorithmic game theory to formulate the right questions that might shed some light on this problem. We are hopeful this might lead to some interesting lines of research.

4.5 Metric Space Generalizations

The urinal problem is just one of many situations where physical privacy might be desirable, in a place where people are entering (and leaving) over time. The unfriendly theatre seating arrangement problem described above is one such example (as is any situation with open seating). Another might be called the beach blanket problem: Frankie and Annette arrive at the beach early in the morning and want to decide where to place their beach blanket to allow for the most privacy throughout the day in order to engage in whatever activities they have planned for the day. Under various assumptions about the behavior of later arrivals, what place should they choose? We note that some old fashioned urinals consist of one contiguous urinal with a single central drain. This would correspond to the continuous version of our original urinal problem.

² The designers of men's rooms do make estimates of such quantities in order to decide how many urinals are required for a given site. As one might expect this will depend on the size of a building, what it is used for (e.g., office building, cinema, shopping mall, etc.) and the expected mix of men versus women versus children. For tables of the recommended number of fixtures required see [8].

A natural generalization of the urinal problem that captures all of these might be termed the *metric space privacy problem*. You are given a metric space (M, d) , an ϵ (the *radius of privacy*), and a deterministic or randomized behavior describing the choices of points in the space of later arrivals. Given a configuration (a subset of points in M) choose a point that maximizes the (expected) time until your privacy radius is violated, i.e., a later arrival chooses a point within distance ϵ of your spot. In the theatre seating problem we take M to be the vertices of an $m \times n$ grid with edge weights equal to 1 and $\epsilon = 1$. In the beach blanket problem M may be modelled by a simple polygon (perhaps even a rectangle) with the Euclidean distance measure and ϵ may depend upon how much privacy you need. In the continuous urinal problem M is a unit interval and ϵ is a function of a man's width and privacy needs.

One suspects that some instances of these problems will turn out to be *NP*-hard as it would appear that in some cases you would be forced to solve a version of the obnoxious facility location problem that in some cases is known to be *NP*-hard. Consideration of a given metric space can be combined with dynamics and/or game theory to yield more problems. We feel there is the potential for many interesting open questions in this area.

5 Conclusions

Our main conclusion is that when faced with the decision of what urinal to choose upon entering the men's room, in order to maximize your privacy, you should probably choose the one furthest from the door if it is available and the one next to it is unoccupied. For a vast majority of the (what we consider) natural behaviors that men choosing urinals might follow, this choice is optimal (in the sense defined above). Beyond this observation, we feel that this problem leads to many interesting variations that are worthy of investigating further and we encourage everyone to do more of their thinking while using public restrooms.

Acknowledgments

We thank Pat Morin for pointing out reference [2] and the anonymous referees for many helpful and amusing suggestions. This research was supported in part by Natural Sciences and Engineering Research Council of Canada (NSERC) and Mathematics of Information Technology and Complex Systems (MITACS).

References

1. American Restroom Association, Public Restroom Design Issues, <http://www.americanrestroom.org/design/index.htm>
2. Decker, D.: A Woman's Guide on How to Pee Standing Up, <http://web.archive.org/web/20031202020806/http://www.restrooms.org/standing.html>

3. Flajolet, P.: A Seating Arrangement Problem, <http://algo.inria.fr/libraries/autocomb/fatmen-html/fatmen1.html>
4. Freedman, D., Shepp, L.: An Unfriendly Seating Arrangement Problem. *SIAM Review* 4, 150 (1962)
5. Friedman, H.D., Rothman, D.: Solution to An Unfriendly Seating Arrangement Problem. *SIAM Review* 6, 180–182 (1964)
6. Georgiou, K., Kranakis, E., Krizanc, D.: Random Maximal Independent Sets and the Unfriendly Theater Arrangement Problem. *Discrete Mathematics* 309, 5120–5129 (2009)
7. Hoffman, L.: Computers and Privacy: A Survey. *Computing Surveys* 1, 85–103 (1969)
8. International Code Council, *International Plumbing Code* (2009)
9. Keane, M., Kranakis, E., Krizanc, D., Narayanan, L.: Routing on Delay Tolerant Sensor Networks. In: *Algosensors 2009*, pp. 155–166 (2009)
10. Kleinrock, L.: *Queueing Systems, Volume 1: Theory*. Wiley, Hoboken (1975)
11. Knuth, D.: The Toilet Paper Problem. *American Math. Monthly* 91, 365–370 (1984)
12. MacKenzie, J.K.: Sequential Filling of a Line by Intervals Placed at Random and Its Application to Linear Absorption. *Journal of Chemical Physics* 37, 723–728 (1962)
13. Paola, C.: *A Survey of Obnoxious Facility Location Problems*. TR-99-11, University of Pisa, Dept. of Informatics (1999)
14. Ross, S.M.: *Stochastic Processes*, 2nd edn. John Wiley & Sons, Inc., Chichester (1996)
15. Soifer, S.: The Evolution of the Bathroom and the Implications for Paruresis, <http://paruresis.org/evolution.pdf>
16. Vogt, R., Nascimento, M.A., Harns, J.: On the Tradeoff between User-Location Privacy and Queried-Location Privacy in Wireless Sensor Networks. In: *AdHoc Now 2009*, pp. 241–254 (2009)